



1904 Third Avenue, Suite 105  
Seattle, Washington 98101  
[www.pscleanair.gov](http://www.pscleanair.gov)

---

# **2022**

## Air Quality

## Data Summary

December 2023

## Table of Contents

<b>Table of Contents .....</b>	<b>ii</b>
<b>List of Figures .....</b>	<b>iii</b>
<b>List of Maps .....</b>	<b>iv</b>
<b>List of Tables .....</b>	<b>iv</b>
<b>Appendix – Data Tables .....</b>	<b>v</b>
<b>Executive Summary .....</b>	<b>1</b>
<b>Monitoring Network .....</b>	<b>2</b>
<b>Air Quality Index .....</b>	<b>7</b>
<b>Particulate Matter .....</b>	<b>9</b>
<b>PM<sub>10</sub> .....</b>	<b>9</b>
<b>PM<sub>2.5</sub> Health and Environmental Effects .....</b>	<b>9</b>
<b>Ultrafine Particulate Matter (UFP) .....</b>	<b>9</b>
<b>PM<sub>2.5</sub>: Federal Reference Method and Continuous Methods .....</b>	<b>10</b>
<b>PM<sub>2.5</sub> Daily Federal Standard and Health Goal .....</b>	<b>10</b>
<b>PM<sub>2.5</sub> Annual Federal Standard .....</b>	<b>18</b>
<b>PM<sub>2.5</sub> Continuous Data and Seasonal Variability .....</b>	<b>22</b>
<b>PM<sub>10</sub>: Annual Standard and Modeled Concentrations .....</b>	<b>23</b>
<b>Particulate Matter – PM<sub>2.5</sub> Speciation and Aethalometers .....</b>	<b>25</b>
<b>Speciation Monitoring and Source Apportionment .....</b>	<b>25</b>
<b>Aethalometer Data .....</b>	<b>26</b>
<b>Ozone .....</b>	<b>28</b>
<b>Nitrogen Dioxide .....</b>	<b>32</b>
<b>Carbon Monoxide .....</b>	<b>34</b>
<b>Sulfur Dioxide .....</b>	<b>35</b>
<b>Lead .....</b>	<b>37</b>
<b>Visibility .....</b>	<b>38</b>
<b>Air Toxics .....</b>	<b>40</b>
<b>Relative ranking based on cancer risk &amp; unit risk factors .....</b>	<b>40</b>
<b>Health effects other than cancer .....</b>	<b>42</b>
<b>Air toxics trends .....</b>	<b>43</b>
<b>Definitions .....</b>	<b>58</b>
<b>General Definitions .....</b>	<b>58</b>

### List of Figures

<b>Figure 1: Days Exceeding the PM<sub>2.5</sub> Health Goal at One or More Monitoring Sites.....</b>	<b>11</b>
<b>Figure 2: Daily PM<sub>2.5</sub> Design Values for King County .....</b>	<b>14</b>
<b>Figure 3: Daily PM<sub>2.5</sub> Design Values for King County with wildfire-impacted days removed .....</b>	<b>14</b>
<b>Figure 4: Daily PM<sub>2.5</sub> Design Values for Kitsap County.....</b>	<b>15</b>
<b>Figure 5: Daily PM<sub>2.5</sub> Design Values for Kitsap County with wildfire-impacted days removed.....</b>	<b>15</b>
<b>Figure 6: Daily PM<sub>2.5</sub> Design Values for Pierce County .....</b>	<b>16</b>
<b>Figure 7: Daily PM<sub>2.5</sub> Design Values for Pierce County with wildfire-impacted days removed.....</b>	<b>16</b>
<b>Figure 8: Daily PM<sub>2.5</sub> Design Values for Snohomish County .....</b>	<b>17</b>
<b>Figure 9: Daily PM<sub>2.5</sub> Design Values for Snohomish County with wildfire-impacted days removed..</b>	<b>17</b>
<b>Figure 10: Annual PM<sub>2.5</sub> Design Values for King County.....</b>	<b>18</b>
<b>Figure 11: Annual PM<sub>2.5</sub> Design Values for King County with wildfire-impacted days removed.....</b>	<b>19</b>
<b>Figure 12: Annual PM<sub>2.5</sub> Design Values for Kitsap County.....</b>	<b>19</b>
<b>Figure 13: Annual PM<sub>2.5</sub> Design Values for Kitsap County with wildfire-impacted days removed.....</b>	<b>20</b>
<b>Figure 14: Annual PM<sub>2.5</sub> Design Values for Pierce County .....</b>	<b>20</b>
<b>Figure 15: Annual PM<sub>2.5</sub> Design Values for Pierce County with wildfire-impacted days removed.....</b>	<b>21</b>
<b>Figure 16: Annual PM<sub>2.5</sub> Design Values for Snohomish County .....</b>	<b>21</b>
<b>Figure 17: Annual PM<sub>2.5</sub> Design Values for Snohomish County with wildfire-impacted days removed .....</b>	<b>22</b>
<b>Figure 18: Annual PM<sub>2.5</sub> Black Carbon .....</b>	<b>27</b>
<b>Figure 19: Ozone for Puget Sound Region.....</b>	<b>31</b>
<b>Figure 20: Ozone for Puget Sound Region with wildfire impacted days removed.....</b>	<b>31</b>
<b>Figure 21: Nitrogen Dioxide (NO<sub>2</sub>) (1998–2005; 2017–2022) and Reactive Nitrogen (NO<sub>y</sub> – NO) (2007–2022) for the Puget Sound Region.....</b>	<b>33</b>
<b>Figure 22: Sulfur Dioxide (SO<sub>2</sub>) 1-Hour Maximum Concentrations (3-Year Average of the 99th Percentile) for the Puget Sound Region.....</b>	<b>36</b>
<b>Figure 23: Puget Sound Visibility .....</b>	<b>39</b>
<b>Figure 24: County-wide Visibility.....</b>	<b>39</b>
<b>Figure 25: Ethylene Oxide Annual Average Potential Cancer Risk at Beacon Hill, 2019–2022 .....</b>	<b>44</b>
<b>Figure 26: Formaldehyde Annual Average Potential Cancer Risk at Beacon Hill, 2000–2022 .....</b>	<b>45</b>

<b>Figure 27: Hexavalent Chromium Annual Average Potential Cancer Risk at Beacon Hill, 2005–2013</b>	46
<b>Figure 28: Benzene Annual Average Potential Cancer Risk at Beacon Hill, 2000–2022</b>	47
<b>Figure 29: Arsenic Annual Average Potential Cancer Risk at Beacon Hill, 2003–2022</b>	48
<b>Figure 30: Acetaldehyde Annual Average Potential Cancer Risk at Beacon Hill, 2000–2022</b>	49
<b>Figure 31: Carbon Tetrachloride Annual Average Potential Cancer Risk at Beacon Hill, 2000–2022</b>	50
<b>Figure 32: Chloroform Annual Average Potential Cancer Risk at Beacon Hill, 2000–2022</b>	51
<b>Figure 33: Ethylene Dichloride Annual Average Potential Cancer Risk at Beacon Hill, 2014–2022</b>	52
<b>Figure 34: 1,3-Butadiene Annual Average Potential Cancer Risk at Beacon Hill, 2000–2022</b>	53
<b>Figure 35: Naphthalene Annual Average Potential Cancer Risk at Beacon Hill, 2008–2022</b>	54
<b>Figure 36: Tetrachloroethylene Annual Average Potential Cancer Risk at Beacon Hill, 2000–2022</b>	55
<b>Figure 37: Ethylbenzene Annual Average Potential Cancer Risk at Beacon Hill, 2007–2022</b>	56
<b>Figure 38: Cadmium Annual Average Potential Cancer Risk at Beacon Hill, 2003–2022</b>	57

### List of Maps

<b>Map 1: Active Air Quality Monitoring Station Locations in 2022</b>	3
<b>Map 2: The 98th Percentile 3-Year Average Daily PM<sub>2.5</sub> Concentrations for 2022</b>	12
<b>Map 3: Ozone 3-year Average of 4th Highest 8-hr Value for 2022</b>	29

### List of Tables

<b>Table 1: Air Quality Monitoring Network Parameters 2022</b>	4
<b>Table 2: Air Quality Index (AQI) Ratings for 2022</b>	8
<b>Table 3: Five-year DVs for PM<sub>10</sub> concentrations for 2017–2022 – scenario 1</b>	24
<b>Table 4: Five-year DVs for PM<sub>10</sub> concentrations for 2017–2022 – scenario 2</b>	24
<b>Table 5: 2022 Beacon Hill Air Toxics Ranking</b>	42
<b>Table 6: 2022 Calculation and Breakpoints for the Air Quality Index (AQI)</b>	58

### Appendix – Data Tables

<b>Monitoring Methods Used from 1999 to 2022 in the Puget Sound Air shed .....</b>	<b>A-3</b>
<b>Historical Air Quality Monitoring Network .....</b>	<b>A-4</b>
<b>Burn Bans 1988 – 2022 .....</b>	<b>A-9</b>
<b>Particulate Matter (PM2.5) – Federal Reference Method .....</b>	<b>A-11</b>
<b>Particulate Matter (PM2.5) – Federal Equivalent Methods .....</b>	<b>A-12</b>
<b>Particulate Matter (PM2.5) – Continuous – Nephelometer .....</b>	<b>A-13</b>
<b>PM2.5 Speciation Analytes Monitored in 2022 .....</b>	<b>A-14</b>
<b>PM2.5 Black Carbon .....</b>	<b>A-15</b>
<b>Ozone (8-hour concentration) .....</b>	<b>A-16</b>
<b>2022 Beacon Hill Air Toxics Statistical Summary for Air Toxics (units in parts per billion) .....</b>	<b>A-17</b>
<b>2022 Beacon Hill Air Toxics Statistical Summary for Air Toxics (units in nanograms per cubic meter) .....</b>	<b>A-17</b>
<b>Estimates of Air Toxics Risk .....</b>	<b>A-18</b>
<b>2022 Air Toxics Unit Risk Factors .....</b>	<b>A-18</b>
<b>2022 Beacon Hill Potential Cancer Risk Estimates per 1,000,000 – 95th Percentile .....</b>	<b>A-19</b>
<b>2022 Non-cancer Reference Concentrations (RfC) and Hazard Indices &gt;1 .....</b>	<b>A-20</b>
<b>2000-2022 Air Toxics Trends Statistical Summary .....</b>	<b>A-21</b>
<b>Air Quality Standards and Health Goals .....</b>	<b>A-22</b>



## 2022 Air Quality Data Summary

The 2022 Air Quality Data Summary is available for viewing or download on the internet at:

**[www.pscleanair.gov](http://www.pscleanair.gov)**

Links to additional documents for download are also available at the web site.



This material is available in alternate formats for people with disabilities.  
Please call Joanna Cruse at 206-689-4067

### Executive Summary

The Puget Sound Clean Air Agency (the Agency) summarizes air quality data from our core monitoring network every year. This report summarizes regional air quality by presenting air quality monitoring results for six criteria air pollutants and air toxics. The U.S. Environmental Protection Agency (EPA) sets national ambient air quality standards (NAAQS) for the criteria pollutants. The criteria pollutants are:

- Particulate Matter (particles 10 micrometers and smaller [PM<sub>10</sub>] and 2.5 micrometers and smaller in diameter [PM<sub>2.5</sub>])
- Ozone
- Nitrogen Dioxide
- Carbon Monoxide
- Sulfur Dioxide
- Lead (monitoring discontinued due to very low levels)

Air toxics are defined by Washington State and the Agency to include hundreds of chemicals and compounds that are associated with a broad range of adverse health effects, including cancer.<sup>1</sup> Many air toxics are a component of either particulate matter or volatile organic compounds (a precursor to ozone). The Air Quality Index (AQI)<sup>2</sup> is a nationwide reporting standard for the criteria pollutants. The AQI is used to relate air quality levels to health effects in a simplified way and is intended mainly for forecasting and real-time communication. “Good” AQI days continued to be in the majority 2022. However, air quality degraded into “moderate” for over 100 days in King and Snohomish Counties, and “unhealthy for sensitive groups”, “unhealthy”, and “very unhealthy” for brief periods. There was one hazardous day in Snohomish County in 2022.

The Agency and the Washington State Department of Ecology (Ecology) work together to monitor air quality within the Puget Sound region. The Agency’s jurisdiction includes King, Kitsap, Pierce, and Snohomish Counties. Real-time air monitoring data are available for pollutants at [www.pscleanair.gov/157/Request-Air-Quality-Data](http://www.pscleanair.gov/157/Request-Air-Quality-Data).

To receive the Agency’s most updated news and stay current on air quality issues in King, Kitsap, Pierce and Snohomish counties, visit [www.pscleanair.gov/258/Connect-With-Us](http://www.pscleanair.gov/258/Connect-With-Us) and select your favorite news feed method. Friends and subscribers receive the latest on air quality news and updates on projects in the Puget Sound region. You can also find us on Facebook and Twitter.

Data included in this report are for our core monitoring network. We also perform local, seasonal monitoring studies – you can see reports on these study results at the library on our website at [www.pscleanair.gov](http://www.pscleanair.gov).

<sup>1</sup>Washington Administrative Code 173-460. See Table of Toxic Air Pollutants, WAC 173-460-150.

[apps.leg.wa.gov/WAC/default.aspx?cite=173-460-150](http://apps.leg.wa.gov/WAC/default.aspx?cite=173-460-150)

<sup>2</sup> <https://www.airnow.gov/aqi/aqi-basics/>

Over the last two decades, many pollutant levels have declined, and air quality has improved overall. In 2022, the overall air quality remained good, continuing the trend of improvement, though we still face challenges due to wildfire smoke. Elevated fine particle levels ( $PM_{2.5}$ ) pose the greatest air quality challenge in our jurisdiction. While fine particle levels met the U.S. Environmental Protection Agency's (EPA's) health-based standard of 35 micrograms per cubic meter in 2022 when days with wildfire smoke are excluded, the Agency's more stringent local  $PM_{2.5}$  health goal of 25 micrograms per cubic meter was exceeded on 40 days which were during wildfire smoke (24 days) and winter months (15 days) at various sites.

Ozone levels remain a concern in our region, however it has shown improvement as values have reduced at various sites. In 2022, all sites except for Enumclaw met the U.S. Environmental Protection Agency's (EPA's) revised 2015 standard. Enumclaw was at the standard of 0.07 ppm.

Some air toxics were measured at levels known to cause adverse health effects. These health effects include, but are not limited to, increased cancer risk, respiratory effects, and developmental effects.


Overall, the air quality in Puget Sound region has continued meeting the standards. However, in September and October 2022, we had various wildfires impacting the air quality at our monitoring sites for six and nineteen days, respectively. Due to the vicinity of some of the wildfires in the region (like Bolt Creek and White River fires), we observed air quality levels reaching very unhealthy and hazardous categories in King and Snohomish counties. Also, the duration of wildfire smoke impacted days were unprecedented, and most of October had levels MODERATE or above across the region.

### Monitoring Network

The Puget Sound Clean Air Agency (the Agency) and the Washington State Department of Ecology (Ecology) operated the monitoring network within the Agency's jurisdiction of King, Kitsap, Pierce, and Snohomish Counties in 2022. The network is comprised of meteorological and pollutant-specific monitors, as well as instruments dedicated to special studies. Data from the network are normally collected automatically via Ecology's data network, or in some cases, collected manually by field staff. Monitoring stations are located in a variety of geographic locations in the Puget Sound region. Monitors are sited according to EPA criteria to ensure a consistent and representative picture of air quality.

Map 1 and Table 1 show King, Kitsap, Pierce, and Snohomish County monitoring sites used in 2022. An interactive map is available at [www.pscleanair.gov/NetworkMap](http://www.pscleanair.gov/NetworkMap).

Map 1: Active Air Quality Monitoring Station Locations in 2022





PUGET SOUND

Clean Air Agency

## 2022 Air Quality Data Summary

**Table 1: Air Quality Monitoring Network Parameters 2022**

Station ID	Location	PM <sub>2.5</sub>					O <sub>3</sub>	SO <sub>2</sub>	NO <sub>x</sub>	CO	b <sub>sp</sub>	Wind	Temp	AT	Vsby	Location	
		Ref	Spec	FEM	Is	bc											
BK◎	10 <sup>th</sup> & Weller, Seattle SPECIATION SITE		●	●		●			●	●			●	●	●		a
BL	11675 44 <sup>th</sup> Ave S, Tukwila Allentown			●	●	●						●	●	●		●	b, e, f
BW◎	Beacon Hill, 15th S & Charlestown, Seattle SPECIATION SITE	●	●	●				●	●	●	●		●	●	●		b, d, f
CE	Duwamish, 4700 E Marginal Way S, Seattle SPECIATION SITE	●	●	●	●	●						●	●	●	●	●	a, e
CW	James St & Central Ave, Kent			●	●	●						●	●	●		●	b, d
DB	17171 Bothell Way NE, Lake Forest Park				●							●				●	b, d, f
DD	South Park, 8201 10 <sup>th</sup> Ave S, Seattle				●							●				●	b, e, f
DF◎	30525 SE Mud Mountain Road, Enumclaw						●						●	●			c
DG◎	42404 SE North Bend Way, North Bend				●		●					●		●		●	c, d, f
DN◎	20050 SE 56 <sup>th</sup> , Lake Sammamish State Park, Issaquah						●										b, d
EQ	Tacoma Tideflats, 2301 Alexander Ave, SPECIATION SITE		●	●	●	●						●	●	●	●	●	a, e
ES	7802 South L St, Tacoma SPECIATION SITE	●	●	●	●	●						●	●	●	●	●	b, f
FG◎	Mt Rainier National Park, Jackson Visitor Center						●										c
IG	Marysville JHS, 1605 7 <sup>th</sup> St, Marysville			●	●							●	●	●		●	b, d
IK	14310 SE 12 <sup>th</sup> St, Bellevue				●							●				●	a, d
JO	Darrington High School, Darrington 1085 Fir St			●	●							●	●	●		●	d, f
PA	1802 S 36th St, Tacoma			●		●				●			●	●	●		a, f



PUGET SOUND

Clean Air Agency

## 2022 Air Quality Data Summary

Station ID	Location	PM <sub>2.5</sub>					O <sub>3</sub>	SO <sub>2</sub>	NO <sub>y</sub>	CO	b <sub>sp</sub>	Wind	Temp	AT	Vsby	Location
		Ref	Spec	FEM	Is	bc										
QK	Spruce, 3250 Spruce Ave, Bremerton			●	●						●	●	●		●	b, f
RV	Yelm N Pacific Road, 931 Northern Pacific Rd SE, Yelm							●								c, f
ZB	Auburn 29th Street, 402 29th Street, Auburn			●	●						●	●	●		●	b

◎	Station operated by Ecology	SO <sub>2</sub>	Sulfur Dioxide
●	Indicates parameter currently monitored	NO <sub>y</sub>	Nitrogen Oxides
PM <sub>2.5</sub> ref	Particulate matter <2.5 micrometers (reference)	CO	Carbon Monoxide
PM <sub>2.5</sub> Spec	Speciation	b <sub>sp</sub>	Light scattering by atmospheric particles (nephelometer)
PM <sub>2.5</sub> FEM	Particulate matter <2.5 micrometers (TEOM-fdms continuous or beta attenuation continuous)	Wind	Wind direction and speed
PM <sub>2.5</sub> Is	Particulate matter <2.5 micrometers (light scattering nephelometer continuous)	Temp	Air temperature (relative humidity also measured at BW, IG, ES)
PM <sub>2.5</sub> bc	Particulate matter <2.5 micrometers black carbon (light absorption aethalometer)	AT	Air Toxics
O <sub>3</sub>	Ozone (May through September except Beacon Hill and Mt Rainier)	VSBY	Visual range (light scattering by atmospheric particles)
Location			
a	Urban Center	d	Commercial
b	Suburban	e	Industrial
c	Rural	f	Residential

## 2022 Air Quality Data Summary

Page A-3 of the Appendix shows a list of the methods used for monitoring the criteria pollutants. Additional information on these methods is available at EPA's website at [epa.gov/ttn/amtic/](http://epa.gov/ttn/amtic/). Information on air toxics monitoring methods is available at [epa.gov/ttn/amtic/airtox.html](http://epa.gov/ttn/amtic/airtox.html).

The Agency has been conducting air quality monitoring as early as 1965. A summary of the monitoring stations and parameters used over the history of the program is on page A-4 of the Appendix. The network changes periodically because the Agency and Ecology regularly re-evaluate monitoring objectives, resources, and logistics. You can also find a list of historical data summaries going back to 1972 here: <http://dl.pscleanair.org/Datasummaries/>.

### Air Quality Index

EPA established the air quality index (AQI) as a simplified tool for communicating daily air quality forecasts and near real-time information. It is intended to help people plan their daily activities. The AQI indicates how clean or polluted air is and what associated health effects might be a concern. It focuses on health effects that may be experienced within a few hours or days after breathing polluted air. EPA calculates the AQI for five major air pollutants regulated by the Clean Air Act: ground-level ozone, particle pollution (also known as particulate matter or PM), carbon monoxide, sulfur dioxide, and nitrogen dioxide.

EPA mainly developed the AQI as a daily indicator or forecast of air quality. To view the real-time AQI for your area, visit <http://www.airnow.gov>. For more information about local air quality, visit [www.pscleanair.gov/27/Air-Quality](http://www.pscleanair.gov/27/Air-Quality).

A higher AQI indicates higher levels of air pollution and greater health concern. An AQI value of 100 generally corresponds to the national air quality standard for the pollutant, which is the level EPA has set to protect public health. It's important to note that health effects can be experienced even at "good" or "moderate" levels.

The purpose of the AQI is to help people understand what local air quality means to health. To make it easier to understand, the AQI is divided into six categories:

Air Quality Index (AQI) Values	Levels of Health Concern	Colors
When the AQI is:	...air quality condition is:	...look for this color:
<b>0 – 50</b>	<b>Good</b>	<b>Green</b>
<b>51 – 100</b>	<b>Moderate</b>	<b>Yellow</b>
<b>101 – 150</b>	<b>Unhealthy for Sensitive Groups</b>	<b>Orange</b>
<b>151 – 200</b>	<b>Unhealthy</b>	<b>Red</b>
<b>201 – 300</b>	<b>Very Unhealthy</b>	<b>Purple</b>
<b>301 – 500</b>	<b>Hazardous</b>	<b>Maroon</b>

Table 1Table 2 shows the percentage of days in each AQI category by county for 2022. The maximum AQI value from all of our network monitors in a county determines its AQI category for the day. Most days were in the "Good" air quality category with some "Moderate" days, some "Unhealthy for

## 2022 Air Quality Data Summary

Sensitive Groups" days, seven wildfire-impacted days that were "Unhealthy", three wildfire-impacted days that was "Very Unhealthy", and one wildfire-impacted day that was "Hazardous".

**Table 2: Air Quality Index (AQI) Ratings for 2022**

County	AQI Rating (% of year)					Highest AQI
	Good	Moderate	Unhealthy for Sensitive Groups	Unhealthy	Very Unhealthy	
King	71.8%	23.3%	3.6%	1.1%	0.3%	212
Kitsap	91.5%	7.7%	0.8%	0.0%	0.0%	143
Pierce	77.0%	20.3%	2.2%	0.5%	0.0%	171
Snohomish	70.1%	23.6%	3.6%	1.6%	0.8%	306



### Particulate Matter

"Particulate matter," also known as particle pollution or PM, is a complex mixture of extremely small particles and liquid droplets. Particle pollution consists of several components, including acids (such as nitrates and sulfates), organic chemicals, metals, and soil or dust particles. PM can be categorized into three broad classes based on size: Coarse—with a diameter of 10  $\mu\text{m}$  or less ( $\text{PM}_{10}$ ); Fine—with a diameter of 2.5  $\mu\text{m}$  or less ( $\text{PM}_{2.5}$ ) and Ultrafine—with a diameter of less than 0.1  $\mu\text{m}$  (UFP).

#### **PM<sub>10</sub>**

$\text{PM}_{10}$  is particulate matter with a diameter of 10 micrometers (or microns) or less. These particles can include larger particles like dust, and smaller particles ( $\text{PM}_{2.5}$ ) that come mainly from combustion sources. Studies show that the finer  $\text{PM}_{2.5}$  particles have more significant health risks. With levels well below the federal standard for years, the Agency ceased direct  $\text{PM}_{10}$  monitoring in 2006. For a historic look at the  $\text{PM}_{10}$  levels in the Puget Sound Region, please request a copy of the 2007 data summary, pages 32–35.<sup>3</sup>

#### **PM<sub>2.5</sub> Health and Environmental Effects**

$\text{PM}_{2.5}$  (or fine particulate matter) has a diameter of 2.5 microns or less. An extensive body of scientific evidence shows that exposure to particle pollution is linked to a variety of significant health problems, such as increased hospital admissions and emergency department visits for cardiovascular and respiratory problems, heart attacks and premature death. Older adults, children, pregnant women, and those with pre-existing health conditions are more at risk from exposure to particle pollution. Particle pollution also reduces visibility in cities and some of our nation's most treasured national parks.

Fine particles are emitted directly from a variety of sources, including wood burning (both outside, and in wood stoves and fireplaces), vehicles and industry. They also form when gases from some of these same sources react in the atmosphere.

#### **Ultrafine Particulate Matter (UFP)**

Emerging health studies indicate that very tiny ultrafine particles with a diameter of 0.1 micron and less may be linked with negative health effects. Currently, there are no health-based standards on what a healthy level of ultrafine particles is. We are exploring new methods for measuring and assessing ultrafine particles, but this technology is not yet ready to add to our core monitoring network.

---

<sup>3</sup> To request a document, please visit [www.pscleanair.gov/272/2396/Records-Request](http://www.pscleanair.gov/272/2396/Records-Request)

### PM<sub>2.5</sub>: Federal Reference Method and Continuous Methods

Fine particulate matter (PM<sub>2.5</sub>) is measured using a variety of methods to ensure quality and consistency. EPA has defined a filter-based method as the federal reference method (FRM)—the primary method used to determine PM<sub>2.5</sub> concentrations. EPA further defined several federal equivalent methods (FEM), which are continuous instruments operated under specific standard operating procedures. The main advantage of continuous FEMs is to provide PM concentrations at a higher temporal resolution (hourly averages) compared to the FRM (24-hour averages).

The Agency uses the FRM, FEMs, and a nephelometer estimation method to provide data. These methods determine fine particulate matter concentrations differently:

- The FRM involves pulling in air (at a given flow rate) for a 24-hour period and collecting particles with a diameter of 2.5 microns or smaller on a filter. The filter is weighed, and the mass is divided by air volume (determined from flow rate and amount of time) to provide concentration. Particles on the filter can later be analyzed for more information about the types of particulate matter.
- We operate a FEM instrument used in the network: the Met-One BAM, a beta attenuation monitor which uses the attenuation of beta radiation to assess the PM<sub>2.5</sub> mass on a filter tape.
- The nephelometer measures the scattering of light in a photomultiplier tube; its results are then compared to FRM and FEM method data to produce an estimate of PM<sub>2.5</sub>. While light scattering has been proven to correlate well with direct PM<sub>2.5</sub> measurements, this is an “unofficial” method because it does not measure particle mass directly.


The Agency and Ecology work together on quality assurance to ensure the FEM-generated data are directly comparable to those generated by the reference method.

### PM<sub>2.5</sub> Daily Federal Standard and Health Goal


The EPA set a daily health-based fine particle standard of 35 micrograms per cubic meter ( $\mu\text{g}/\text{m}^3$ ). Monitors in King, Kitsap, Pierce, and Snohomish Counties exceeded  $35 \mu\text{g}/\text{m}^3$  26 days in 2022, 17 of which were due to wildfire smoke. The majority of the remaining exceedance days were in November and December. In addition to the federal standard, our Board of Directors adopted a more stringent health goal of  $25 \mu\text{g}/\text{m}^3$  in 1999, based on recommendations from our Particulate Matter Health Committee. Monitors in our four counties exceeded the local health goal of  $25 \mu\text{g}/\text{m}^3$  on 40 days which were mostly during wildfire-impacted days in September and October, and in the winter months in 2022.

Figure 1 shows the number of days the health goal was exceeded annually in the region, from 2000 to 2022. Our region exceeded the health goal on 33 days in the heating season and 7 days in the non-heating seasons as well. A majority of these exceedance days were due to wildfire smoke in September and October.

**Figure 1: Days Exceeding the PM<sub>2.5</sub> Health Goal at One or More Monitoring Sites**



Map 2 shows the 98<sup>th</sup> percentile of the 3-year average of daily PM<sub>2.5</sub> concentrations from 2020 to 2022. This map incorporates data collected from federal reference, federal equivalent, and nephelometer estimation methods. The map also does not show sites with less than three years or incomplete data.

Map 2: The 98th Percentile 3-Year Average Daily PM<sub>2.5</sub> Concentrations for 2022



PUGET SOUND


Clean Air Agency

## 2022 Air Quality Data Summary

Figures 2 through 9 show daily 98<sup>th</sup> percentile 3-year averages at each monitoring station in King, Kitsap, Pierce, and Snohomish Counties compared to the current daily federal standard. Points on the graphs represent averages for three consecutive years. For example, the value for 2022 is the average of the 98<sup>th</sup> percentile daily concentration for 2020, 2021, and 2022. These figures incorporate data collected from federal reference, federal equivalent, and nephelometer estimation methods. Due to Covid-19, federal reference monitoring could not be conducted from mid-March to July 2020, hence for these months only federal equivalent and nephelometer monitoring data has been used.


For each county, we include two figures: the first shows the entire dataset, and the second shows levels with wildfire smoke-impacted days removed in 2017, 2018, 2020, 2021, and 2022. The EPA allows data from days that were influenced by exceptional events that are beyond the ability of air agencies to control, such as wildfires or dust storms, to be excluded from regulatory calculations. There were 26 wildfire-impacted days in September and October 2022. With wildfire smoke-impacted days excluded from 2020, 2021, and 2022, all monitors in our four counties fall below the federal standard of 35  $\mu\text{g}/\text{m}^3$ . Without excluding wildfire smoke-impacted days, monitors in King and Snohomish Counties equaled or exceeded the standard in 2022.

Figures 4 and 5 do not show any 2012–2014 data for Kitsap County because the Bremerton monitoring site moved to a new location and design values could not be computed until three complete years of data were collected at the new site. Statistical summaries for 98<sup>th</sup> percentile daily concentrations for 2022 data are provided on pages A-11 through A-13 of the Appendix.

Figure 2: Daily PM<sub>2.5</sub> Design Values for King County


Note: Duwamish data are FRM from 1999–2005, 2007–09, nephelometer 2006, 2010, FEM 2011–2020. Beacon Hill data are FRM from 1999–2009, FEM 2010–20. Lake Forest Park data are FRM from 1999–2007, nephelometer in 2008–2020. South Park data are FRM from 1999–2004, nephelometer in 2005–2020. Bellevue Way data are FRM from 2001–2004, nephelometer 2005–20. Redmond data are FRM from 2000–2002, nephelometer from 2003–2005. Queen Anne data are nephelometer from 2002–2015. Olive Way data are nephelometer from 2003–2013. North Bend data are FRM from 2000–2004, nephelometer in 2005–2020. Kent data are FRM from 1999–2004, nephelometer in 2005–2010, FEM 2011–2020. Enumclaw data are from nephelometer in 2000–2009.


\* Indicates an estimate based on incomplete data. Data less than 75% in a quarter at South Park in 2002 & 2016, Beacon Hill in 2008, Bellevue way in 2013, Seattle 10th and Weller in 2022.

 Figure 3: Daily PM<sub>2.5</sub> Design Values for King County with wildfire-impacted days removed


Note: Duwamish data are FRM from 1999–2005, 2007–09, nephelometer 2006, 2010, FEM 2011–2020. Beacon Hill data are FRM from 1999–2009, FEM 2010–20. Lake Forest Park data are FRM from 1999–2007, nephelometer in 2008–2020. South Park data are FRM from 1999–2004, nephelometer in 2005–2020. Bellevue Way data are FRM from 2001–2004, nephelometer 2005–20. Redmond data are FRM from 2000–2002, neph from 2003–2005. Queen Anne data are nephelometer from 2002–2015. Olive Way data are nephelometer from 2003–2013. North Bend data are FRM from 2000–2004, nephelometer in 2005–2020. Kent data are FRM from 1999–2004, nephelometer in 2005–2010, FEM 2011–2020. Enumclaw data are from nephelometer in 2000–2009.

\* Indicates an estimate based on incomplete data. Data less than 75% in a quarter at South Park in 2002 & 2016, Beacon Hill in 2008, Bellevue way in 2013, Seattle 10th and Weller in 2022.


**Figure 4: Daily PM<sub>2.5</sub> Design Values for Kitsap County**



Note: Bremerton Meadowdale site ended 4/30/12 and Bremerton Spruce site began 5/1/2012. Bremerton Meadowdale data are nephelometer from 2006-12. Bremerton Spruce data are FEM from 2012-20.

\* Indicates an estimate based on incomplete data. Data less than 75% in two quarters at Bremerton Meadowdale in 2008.

**Figure 5: Daily PM<sub>2.5</sub> Design Values for Kitsap County with wildfire-impacted days removed**




Note: Bremerton Meadowdale site ended 4/30/12 and Bremerton Spruce site began 5/1/2012. Bremerton Meadowdale data are neph from 2006-12. Bremerton Spruce data are FEM from 2012-20.

\* Indicates an estimate based on incomplete data. Data less than 75% in two quarters at Bremerton Meadowdale in 2008.




**Figure 6: Daily PM<sub>2.5</sub> Design Values for Pierce County**



Note: South L data are FRM from 2000–2020. Tacoma Tideflats data are FRM from 1999–2002 and nephelometer from 2003–2020. Puyallup data are FRM from 1999–2002 and nephelometer from 2003–2004 and 2006–2019. Puyallup site has been discontinued in 2020.


\* Indicates an estimate based on incomplete data. Data less than 75% in two quarters at Puyallup in 2005.

**Figure 7: Daily PM<sub>2.5</sub> Design Values for Pierce County with wildfire-impacted days removed**




Note: South L data are FRM from 2000–2020. Alexander Avenue data are FRM from 1999–2002 and nephelometer from 2003–2020. Puyallup data are FRM from 1999–2002 and nephelometer from 2003–2004 and 2006–2019. Puyallup site has been discontinued in 2020.

\* Indicates an estimate based on incomplete data. Data less than 75% in two quarters at Puyallup in 2005.

Figure 8: Daily PM<sub>2.5</sub> Design Values for Snohomish County


Note: Marysville data are FRM 1999–2011, FEM 2012–20. Lynnwood data are FRM 2002–03, 2005–06, Nephelometer 2004, 2007–10, FEM 2011–2016, FEM & Nephelometer combined in 2017, Nephelometer 2018–19. Darrington data are nephelometer in 2006, FRM in 2007–2011, FEM 2012–20. Lynnwood site has been discontinued in 2020.

\* Indicates an estimate based on incomplete data. Data less than 75% for a quarter at Marysville in 2017

 Figure 9: Daily PM<sub>2.5</sub> Design Values for Snohomish County with wildfire-impacted days removed


Note: Marysville data are FRM 1999–2011, FEM 2012–20. Lynnwood data are FRM 2002–03, 2005–06, Neph 2004, 2007–10, FEM 2011–2016, FEM & Neph combined in 2017, Neph 2018–19. Darrington data are neph in 2006, FRM in 2007–2011, FEM 2012–20. Lynnwood site has been discontinued in 2020.

\* Indicates an estimate based on incomplete data. Data less than 75% for a quarter at Marysville in 2017.

**PM<sub>2.5</sub> Annual Federal Standard**


Figures 10 through 17 present 3-year average of annual concentrations at each monitoring station for King, Kitsap, Pierce, and Snohomish Counties. In 2012, the EPA strengthened the annual standard from 15  $\mu\text{g}/\text{m}^3$  to 12  $\mu\text{g}/\text{m}^3$ . All counties have levels below the 12  $\mu\text{g}/\text{m}^3$  annual standard. Figures 12 and 13 do not show any 2012–2014 data for Kitsap County because the Bremerton monitoring site moved to a new location and design values could not be computed until three complete years of data were collected at the new site.

Figures 10 through 17 include data from the federal reference method (FRM) and continuous method monitors. The federal standard is based on a 3-year average, and each value on the graph is an average of the current year and the two prior years. For example, the value shown for 2022 is the average of the annual averages for 2020, 2021, and 2022. As with the daily standard, for each county we include two figures: the first shows the entire dataset, and the second shows levels with wildfire smoke-impacted days removed in 2017, 2018, 2020, 2021, and 2022.

**Figure 10: Annual PM<sub>2.5</sub> Design Values for King County**

### King County PM<sub>2.5</sub> Annual Design Values


(3-year average of annual mean concentrations)



Note: Duwamish data are FRM from 1999–2005, 2007–09, nephelometer 2006, 2010, FEM 2011–2020. Beacon Hill data are FRM from 1999–2009, FEM 2010–20. Lake Forest Park data are FRM from 1999–2007, nephelometer 2008–2020. South Park data are FRM from 1999–2004, nephelometer 2005–2020. Bellevue Way data are FRM from 2001–2004, nephelometer 2005–20. Redmond data are FRM from 2000–2002, nephelometer 2003–2005. Queen Anne data are nephelometer from 2002–2015. Olive Way data are nephelometer from 2003–2013. North Bend data are FRM from 2000–2004, nephelometer 2005–2020. Kent data are FRM from 1999–2004, nephelometer 2005–2010, FEM 2011–2020. Enumclaw data are nephelometer from 2000–2009.

\*Indicates an estimate based on incomplete data. Data less than 75% complete in one quarter at South Park in 2002 & 2016, Beacon Hill in 2008, Bellevue way in 2013, Seattle 10th and Weller in 2022.


**Figure 11: Annual PM<sub>2.5</sub> Design Values for King County with wildfire-impacted days removed**



Note: Duwamish data are FRM from 1999-2005, 2007-09, nephelometer 2006,2010, FEM 2011-2020. Beacon Hill data are FRM from 1999-2009, FEM 2010-20. Lake Forest Park data are FRM from 1999-2007, nephelometer 2008-2020. South Park data are FRM from 1999-2004, nephelometer 2005-2020. Bellevue Way data are FRM from 2001-2004, nephelometer 2005-20. Redmond data are FRM from 2000-2002, nephelometer 2003-2005. Queen Anne data are nephelometer from 2002-2015. Olive Way data are nephelometer from 2003-2013. North Bend data are FRM from 2000-2004, nephelometer 2005-2020. Kent data are FRM from 1999-2004, nephelometer 2005-2010, FEM 2011-2020. Enumclaw data are nephelometer from 2000-2009.

\*Indicates an estimate based on incomplete data. Data less than 75% complete in one quarter at South Park in 2002 & 2016, Beacon Hill in 2008, Bellevue way in 2013, Seattle 10th and Weller in 2022.

**Figure 12: Annual PM<sub>2.5</sub> Design Values for Kitsap County**




Note: Bremerton Meadowdale site ended 4/30/12 and Bremerton Spruce site began 5/1/2012. Bremerton Meadowdale data are nephelometer from 2006-12. Bremerton Spruce data are FEM from 2012-20.


\* Indicates an estimate based on incomplete data. Data less than 75% complete in two quarters at Bremerton Meadowdale in 2008.



**Figure 13: Annual PM<sub>2.5</sub> Design Values for Kitsap County with wildfire-impacted days removed**




**Figure 14: Annual PM<sub>2.5</sub> Design Values for Pierce County**






**Figure 15: Annual PM<sub>2.5</sub> Design Values for Pierce County with wildfire-impacted days removed**



Note: South L data are FRM from 2000–2020. Tacoma Tideflats data are FRM from 1999–2002 and nephelometer from 2003–2020. Puyallup data are FRM from 1999–2002 and nephelometer from 2003–2004 and 2006–2019. Puyallup site has been discontinued in 2020.

\* Indicates an estimate based on incomplete data. Data less than 75% complete at Puyallup in two quarters in 2005 & one quarter in 2015, one quarter at Tacoma Tideflats in 2015.

**Figure 16: Annual PM<sub>2.5</sub> Design Values for Snohomish County**




Note: Marysville data are FRM 1999–2011, FEM 2012–20. Lynnwood data are FRM 2002–03, 2005–06, Nephelometer 2004, 2007–10, FEM 2011–2016, FEM & Nephelometer combined in 2017, Nephelometer 2018–19. Darrington data are neph in 2006, FRM in 2007–2011, FEM 2012–20. Lynnwood site has been discontinued in 2020.

\* Indicates an estimate based on incomplete data. Data less than 75% complete for a quarter at Marysville in 2017.



**Figure 17: Annual PM<sub>2.5</sub> Design Values for Snohomish County with wildfire-impacted days removed**



### PM<sub>2.5</sub> Continuous Data and Seasonal Variability

Continuous monitoring data provide information on how PM<sub>2.5</sub> levels vary throughout the year. For example, many sites have elevated PM<sub>2.5</sub> levels during the winter when residential wood burning and air stagnations are at their peak but have low levels of PM<sub>2.5</sub> during the summer. A summary of the continuous data for PM<sub>2.5</sub>, black carbon, and ozone for the year 2022 is available at <https://pscleanair.gov/615/Data-Summary>. For more detailed information on continuous data, please see the Air Graphing tool at <https://secure.pscleanair.org/airgraphing> to plot the sites and timeframes of interest.

### PM<sub>10</sub>: Annual Standard and Modeled Concentrations

Our region was nonattainment for PM<sub>10</sub> in 1987 in the three industrial areas of Puget Sound: Seattle Duwamish Valley, Tacoma Tideflats, and Kent. The streets were paved, and the area saw significant reductions thereafter and levels were far below the standard since. Our region completed all its [limited maintenance plan](#) obligations in 2020, including monitoring for PM<sub>10</sub>.

While the direct monitoring of PM<sub>10</sub> concentrations ended in 2007, we can still model recent concentration levels of PM<sub>10</sub> using the observed PM<sub>2.5</sub> concentrations and two site-dependent linear relationships (one for summer (Apr-Sep) and one for winter (Oct-Mar)). These relationships were established for the 1999–2007 time-period when PM<sub>10</sub> and PM<sub>2.5</sub> were recorded simultaneously at our sites. The main assumption with that method is that the linear relationships remained constant over time.<sup>4</sup>

In Table 3 and Table 4, we present the design values (DVs) that have been calculated using both a table-look-up method and a statistical-fit method, described in the EPA PM<sub>10</sub> State Implementation Plan (SIP) Development Guideline<sup>5</sup>. We did these calculations for the following sites: Kent (AQS Site ID: 53 033 2004), Seattle–Duwamish (53 033 0057), and Tacoma Tideflats (53 053 0031) PM<sub>10</sub> Maintenance Areas and for the last five years (2017–2022).

The DVs presented in Table 3 and Table 4 are calculated following two scenarios:

- **Scenario 1 (Table 3):** All daily modeled PM<sub>10</sub> concentrations are included in the DV calculation.
- **Scenario 2 (Table 4):** Daily modeled PM<sub>10</sub> concentrations are excluded from the DV calculation during 2017, 2018, 2020, 2021 & 2022 wildfire-smoke days (I-Flags<sup>6,7,8,9,10</sup>).

Five-year DVs less than 98 µg/m<sup>3</sup> were required to continue to qualify for the Limited Maintenance Plan (LMP)<sup>9</sup>. Over the last three years, scenario 1 (with wildfire smoke days included) would not have met this qualification. Scenario 2 would have met the qualification and is a more appropriate

---

<sup>4</sup> The PM<sub>2.5</sub> concentrations come from several instruments at each site. At all sites, we prioritize instruments measuring PM<sub>2.5</sub> concentrations with missing values in the following way: FEM BAM > 1400ab/8500 FEM TEOM > 1405 FEM TEOM > nephelometer. While Kent and Seattle–Duwamish have most of their data coming from TEOM (2013–2018) and BAM (2018–2021), only a nephelometer has been in operation at Tacoma–Tideflats for 2013–2021. As of May 26, 2021, a BAM has been added a Tacoma–Tideflats and its data will be used next year once a full year of monitoring will be completed.

<sup>5</sup> PM<sub>10</sub> SIP Development Guideline – United States Environmental Protection Agency. June 1987. EPA-450/2-86-001

<sup>6</sup> The Tacoma Tideflats monitoring site experienced higher than typical firework activity in the late June – early July of 2021 leading to July 4<sup>th</sup> with elevated PM<sub>2.5</sub> concentrations explaining the higher 2021 DV.

<sup>7</sup> Informational Flag request for 2017 Wildfire Affected Exceedances – WA Dept. of Ecology. Flagging Memo. Feb 2018.

<sup>8</sup> Informational Flag request for 2018 Wildfire Affected Exceedances – WA Dept. of Ecology. Flagging Memo. Feb 2019.

<sup>9</sup> Informational Flag request for 2020 Wildfire Affected Exceedances – WA Dept. of Ecology. Flagging Memo. May 2021.

<sup>10</sup> Informational Flag request for 2021 Wildfire Affected Exceedances – WA Dept. of Ecology. Flagging Memo. April 2022.

<sup>10</sup> Informational Flag request for 2022 Wildfire Affected Exceedances – WA Dept. of Ecology 2023.



estimate for the Puget Sound region's three former Maintenance Areas, given the unprecedented wildfire smoke levels witnessed in summers of 2017, 2018, 2020, 2021 & 2022.

**Table 3: Five-year DVs for PM<sub>10</sub> concentrations for 2017–2022 – scenario 1**

<b>Sites:</b>		5-yr PM <sub>10</sub> DVs – Scenario 1					
		2017	2018	2019	2020	2021	2022
<b>Kent</b>	89±25 (82)	115±35 (118)	115±37 (118)	226±98 (214)	226±98 (214)	225±104 (214)	
	72±11 (80)	119±45 (110)	117±46 (101)	216±104 (192)	216±103 (192)	217±102 (192)	
<b>Seattle</b>	93±39 (94)	163±69 (165)	163±70 (165)	234±92 (240)	232±90 (240)	229±96 (240)	
<b>Tacoma</b>							

**Table 4: Five-year DVs for PM<sub>10</sub> concentrations for 2017–2022 – scenario 2**

<b>Sites:</b>		5-yr PM <sub>10</sub> DVs – Scenario 2					
		2017	2018	2019	2020	2021	2022
<b>Kent</b>	54±6 (53)	62±12 (65)	60±13 (64)	54±13 (44)	55±12 (56)	51±13 (43)	
	53±3 (56)	52±3 (53)	48±4 (48)	46±3 (46)	46±3 (48)	42±4 (43)	
<b>Seattle</b>	60±16 (55)	60±12 (58)	59±13 (58)	59±14 (57)	73±25 <sup>6</sup> (67)	73±26 <sup>6</sup> (67)	
<b>Tacoma</b>							

Values appear as DV ± an uncertainty interval from a lognormal fit to the data and its 95% prediction interval. Parenthetical values are the DVs obtained using the table-look-up method.<sup>5</sup>

### Particulate Matter – PM<sub>2.5</sub> Speciation and Aethalometers

Although there are no regulatory requirements to go beyond measuring the total mass of fine particulate matter, it is beneficial to know its chemical makeup in addition to its mass. Knowledge about the composition of fine particulate can help guide emissions reduction strategies, such as the Agency's commitment to reducing wood smoke and diesel particulate emissions,<sup>8,9,10</sup> and is useful to scientific and health researchers investigating questions about the effects of fine particulate matter on human health and the environment.

#### Speciation Monitoring and Source Apportionment

Speciation monitoring involves determining the chemical composition of fine particulate matter collected on different types of filters. Speciation filters are analyzed to determine what metals and organic molecules make up the fine particulate at a site. Over 40 chemical species are measured at speciation monitors in the area. These data are used in source apportionment models to estimate contributing sources to PM<sub>2.5</sub>. Source apportionment models use statistical patterns in data to identify likely pollution sources and then estimate how much each source is contributing at each site.

Ecology and PSCAA conducted speciation monitoring at five sites in the Puget Sound region in 2022:

- Seattle Beacon Hill – typical urban impacts, mixture of sources (speciation samples collected every third day, operated by Ecology)
- Seattle 10<sup>th</sup> & Weller – Near-road micro-scale monitoring site (speciation samples collected every sixth day, operated by Ecology)
- Seattle Duwamish – urban site with industrial sources (speciation samples collected every sixth day, operated by PSCAA)
- Tacoma South L – urban residential area, impacts from residential wood combustion (speciation samples collected every sixth day, operated by Ecology)
- Tacoma Tideflats – urban site with industrial sources (speciation samples collected every sixth day, operated by Ecology)

In addition to using speciation data for concentrations of specific species or source apportionment modeling, the Agency uses them to qualitatively look at the makeup of fine particulate at our monitoring sites. For a list of PM<sub>2.5</sub> analytes measured at these sites, please see page A-14 of the Appendix.

---

<sup>8</sup>Puget Sound Air Toxics Evaluation, October 2003. [www.pscleanair.org/DocumentCenter/View/2355/Puget-Sound-Air-Toxics-Evaluation-Final-ReportPDF?bidId=](http://www.pscleanair.org/DocumentCenter/View/2355/Puget-Sound-Air-Toxics-Evaluation-Final-ReportPDF?bidId=)

<sup>9</sup>Tacoma and Seattle Air Toxics Evaluation, October 2010. [epa.gov/ttn/amtic/files/20072008csatam/PSCAA\\_CommunityAssessment\\_FR.pdf](http://epa.gov/ttn/amtic/files/20072008csatam/PSCAA_CommunityAssessment_FR.pdf).

<sup>10</sup>Ogulei, D. WA State Dept of Ecology (2010). "Sources of Fine Particles in the Wapato Hills–Puyallup River Valley PM<sub>2.5</sub> Nonattainment Area". Publication Number 10-02-009. <https://fortress.wa.gov/ecy/publications/documents/1002009.pdf>

### Aethalometer Data


Aethalometers provide information about the carbon fraction of fine particulate matter. Aethalometers continuously measure light absorption at seven different optical wavelengths to estimate carbon concentrations. Two of these wavelengths are important in our evaluation: black carbon (BC) and ultraviolet (UV). Measurements from the black carbon channel correlate well with elemental carbon (EC) concentrations derived from speciation data. Measurements from the UV channel help produce a qualitative estimate of organic carbon (OC), which is correlated with the difference between the UV and BC channel measurements (UV-BC). Elemental and organic carbon are related to diesel particulate, wood smoke particulate, and particulate from other combustion sources.<sup>11</sup> Unfortunately, neither is uniquely attributed to a particular combustion type, so the information gained from aethalometer data is qualitative.

The Agency maintains aethalometers at monitoring sites with high particulate matter concentrations, as well as sites with speciation data, so that data from the different methods to measure carbon may be compared. Figure 18 shows annual average trending of black carbon concentrations. Since 2003, the general trend shows reducing BC levels. A statistical summary of aethalometer black carbon data is presented on page A-15 of the Appendix.

---

<sup>11</sup>Urban Air Monitoring Strategy – Preliminary Results Using Aethalometer Carbon Measurements for the Seattle Metropolitan Area. <https://www3.epa.gov/ttnamtl/archive/files/ambient/samwg/spring2004/awmaurb.pdf>

Figure 18: Annual PM<sub>2.5</sub> Black Carbon



### Ozone

Ozone is a summertime air pollution problem in our region and is not directly emitted by pollutant sources. Ozone forms when photochemical pollutants react with sunlight. These pollutants are called ozone precursors and include volatile organic compounds (VOC) and nitrogen oxides (NO<sub>x</sub>), with some influence by carbon monoxide (CO). These precursors come from human activities such as transportation and solvent use, as well as natural sources. Ozone levels are usually highest in the afternoon because of the intense sunlight and the time required for ozone to form in the atmosphere. The Washington State Department of Ecology conducts ozone monitoring in our four counties.

People sometimes confuse upper atmosphere ozone with ground-level ozone. Upper atmosphere, or stratospheric ozone, helps to protect the earth from the sun's harmful ultraviolet rays. In contrast, ozone formed at ground level is unhealthy. Elevated concentrations of ground-level ozone can cause reduced lung function and respiratory irritation and can aggravate asthma.<sup>12</sup> Ozone has also been linked to immune system impairment. People with respiratory conditions should limit outdoor exertion if ozone levels are elevated. Even healthy individuals may experience respiratory symptoms on a high-ozone day. Ground-level ozone can also damage forests and agricultural crops, interfering with their ability to grow and produce food.<sup>13</sup>

Most ozone monitoring stations are located in rural areas of the Puget Sound region in the western foothills of the Cascade Mountains, while the precursor chemicals that react with sunlight to produce ozone are generated primarily in large metropolitan areas (mostly by cars and trucks). The photochemical formation of ozone takes several hours, and the highest concentrations of ozone are measured in the communities downwind of these large urban areas. In the Puget Sound region, the hot sunny days favorable for ozone formation also tend to have light north-to-northwest winds. Map 3 shows the ozone monitoring network and the design values (three-year average of the 4<sup>th</sup> highest concentrations) measured in 2020–2022.


---

<sup>12</sup>EPA, Air Quality Index: A Guide to Air Quality and Your Health; [epa.gov/airnow/aqi\\_brochure\\_02-14.pdf](https://epa.gov/airnow/aqi_brochure_02-14.pdf).

<sup>13</sup>EPA Health and Environmental Effects of Ground Level Ozone; [epa.gov/ozone-pollution/ozone-basics](https://epa.gov/ozone-pollution/ozone-basics).

## 2022 Air Quality Data Summary

Map 3: Ozone 3-year Average of 4th Highest 8-hr Value for 2022



## 2022 Air Quality Data Summary

Figures 19 and 20 present data for each monitoring station and the 8-hour federal standard. Figure 19 shows levels with the entire dataset, and Figure 20 shows ozone levels with wildfire smoke impacted days removed in 2017, 2018, 2020, 2021, and 2022. The federal standard is based on the 3-year average of the annual 4<sup>th</sup> highest 8-hour concentration, called the “design value”. The year on the x-axis represents the last year averaged. For example, concentrations shown for 2022 are an average of 2020, 2021, and 2022 4<sup>th</sup> highest concentrations.


The EPA’s 2015 8-hour standard is 0.070 ppm. The highest 2022 site design value (for the entire dataset, including wildfire smoke impacted days) is 0.070 ppm at the Enumclaw site. This site would not be considered nonattainment at this time due to the wildfire smoke influence that could be excluded, and an actual exceedance would occur at 0.071 ppm and above. All other monitoring sites are below the standard value of 0.070 ppm.

Statistical summaries for 8-hour average ozone data are provided on page A-16 of the Appendix.


For additional information on ozone, visit <https://www.epa.gov/ozone-pollution>.

## 2022 Air Quality Data Summary

**Figure 19: Ozone for Puget Sound Region**



**Figure 20: Ozone for Puget Sound Region with wildfire impacted days removed**



### Nitrogen Dioxide

Nitrogen dioxide ( $\text{NO}_2$ ) is a reddish brown, highly reactive gas that forms from the reaction of nitrogen oxide (NO) and hydroperoxy ( $\text{HO}_2$ ) and alkylperoxy ( $\text{RO}_2$ ) free radicals in the atmosphere.  $\text{NO}_2$  can cause coughing, wheezing and shortness of breath in people with respiratory diseases such as asthma.<sup>14</sup> Long-term exposure can lead to respiratory infections.

The term  $\text{NO}_x$  is defined as  $\text{NO} + \text{NO}_2$ .  $\text{NO}_x$  participates in a complex chemical cycle with volatile organic compounds (VOCs) which can result in the production of ozone.  $\text{NO}_x$  can also be oxidized to form nitrates, which are an important component of fine particulate matter. On-road vehicles such as trucks and automobiles and off-road vehicles such as construction equipment, marine vessels and port cargo-handling equipment are the major sources of  $\text{NO}_x$  in our region. Industrial boilers and processes, home heaters, and gas stoves also produce  $\text{NO}_x$ .

Motor vehicle and non-road engine manufacturers have been required by EPA to reduce  $\text{NO}_x$  emissions from cars, trucks, and non-road equipment. As a result, emissions have declined dramatically since the 1970s.

EPA promulgated a 1-hour national ambient air quality standard for nitrogen dioxide on January 22, 2010.<sup>15</sup> Since then, Department of Ecology added two “near-road” monitoring sites very close to Interstate 5: one in Seattle (10<sup>th</sup> & Weller), and one in Tacoma (South 36<sup>th</sup> St.). To learn more about the monitoring method visit <https://www.epa.gov/amtic/near-road-monitoring>

In addition to the near-road sites, the Department Ecology measures nitrogen dioxide at the Seattle Beacon Hill site. The monitoring method now records  $\text{NO}_y$  instead of  $\text{NO}_x$ , in order to observe all reactive nitrogen compounds.  $\text{NO}_y$  is  $\text{NO}_x$  plus all other reactive nitrogen oxides present in the atmosphere.  $\text{NO}_y$  components such as nitric acid ( $\text{HNO}_3$ ) and peroxyacetyl nitrate (PAN) can be important contributors to the formation of ozone and fine particulate matter. Direct  $\text{NO}_2$  measurements at Beacon Hill using Cavity Attenuated Phase Shift (CAPS) spectroscopy technique started in 2020.

Figure 21 shows  $\text{NO}_2$  concentrations for Beacon Hill through 2005. In 2006, no data were recorded due to the relocation of the Beacon Hill monitor to a different location on the same property. From 2007 until 2020 at Beacon Hill, the concentration of  $\text{NO}_2$  is represented as  $\text{NO}_y - \text{NO}$ , since  $\text{NO}_2$  was no longer directly recorded, and  $\text{NO}_y = \text{NO} + \text{NO}_2 + \text{other nitroxyl compounds}$ .

<sup>14</sup>EPA, Airnow,  $\text{NO}_x$  Chief Causes for Concern; [epa.gov/airquality/nitrogenoxides/](https://www.epa.gov/airquality/nitrogenoxides/)


<sup>15</sup>EPA. New 1-hour National Ambient Air Quality Standards for Nitrogen Dioxide; [epa.gov/airquality/nitrogenoxides/actions.html](https://www.epa.gov/airquality/nitrogenoxides/actions.html).

## 2022 Air Quality Data Summary

The 2010 1-hour standard is 100 ppb and is based on the 98<sup>th</sup> percentile of 1-hour daily maximum concentrations, averaged over three years. Nitrogen dioxide levels in the Puget Sound region, as currently monitored by Ecology, are typically below (cleaner than) the 1-hour standard. The 1-hour standard is depicted in Figure 21 with historical data since 1998. The years prior to 2010 have been included on the graphs for historical comparison.

Visit [epa.gov/airquality/nitrogenoxides/](https://epa.gov/airquality/nitrogenoxides/) for additional information on NO<sub>2</sub>.

**Figure 21: Nitrogen Dioxide (NO<sub>2</sub>) (1998–2005; 2017–2022) and Reactive Nitrogen (NO<sub>y</sub> – NO) (2007–2022) for the Puget Sound Region**



### Carbon Monoxide

Carbon monoxide (CO) is an odorless, colorless gas that can enter the bloodstream through the lungs and reduce the amount of oxygen that reaches organs and tissues. Carbon monoxide forms when the carbon in fuels does not burn completely. Most of the CO emissions come from motor vehicles.

Elevated levels of CO in ambient air occur more frequently in areas with heavy traffic and during the colder months of the year when temperature inversions are more common. People with cardiovascular disease or respiratory problems may experience chest pain and increased cardiovascular symptoms, particularly while exercising, if CO levels are high. High levels of CO can affect alertness and vision even in healthy individuals.

Although urban portions of the Puget Sound region have historically violated the CO standard, CO levels have decreased significantly primarily due to emissions controls on car engines. EPA designated the Puget Sound region as a CO attainment area in 1996. Ecology has substantially reduced its CO monitoring network, and only the Beacon Hill site remains from the historical network. The near-road site at 10<sup>th</sup> & Weller began operation in June 2014. There currently are no CO monitoring stations in Kitsap, Pierce, or Snohomish Counties.

The CO national ambient air quality standard is based on the 2<sup>nd</sup> highest 8-hour average using the procedures published in the federal register. The EPA also has a 1-hour standard for CO of 35 ppm, not to be exceeded more than once a year. Measured 1-hour concentrations in the Puget Sound area are typically much lower than the 35 ppm standard.

For a historic look at the Puget Sound region's carbon monoxide levels, please see the 2015 Air Quality Data Summary which is available on our website at [www.pscleanair.gov/DocumentCenter/View/2294/Air-Quality-Data-Summary-2015PDF](http://www.pscleanair.gov/DocumentCenter/View/2294/Air-Quality-Data-Summary-2015PDF).

For additional information on CO, visit [epa.gov/airquality/carbonmonoxide](http://epa.gov/airquality/carbonmonoxide).

### Sulfur Dioxide

Sulfur dioxide ( $\text{SO}_2$ ) is a colorless, reactive gas produced by burning fuels containing sulfur, such as coal and oil, and by industrial processes. Historically, the greatest sources of  $\text{SO}_2$  were industrial facilities that derived their products from raw materials such as metallic ore, coal, and crude oil, or that burned coal or oil to produce process heat (petroleum refineries, cement manufacturing and metal processing facilities). Marine vessels, on-road vehicles, and diesel construction equipment are the main contributors to  $\text{SO}_2$  emissions today.

$\text{SO}_2$  may cause people with asthma who are active outdoors to experience bronchial constriction, the symptoms of which include wheezing, shortness of breath and tightening of the chest. People should limit outdoor exertion if  $\text{SO}_2$  levels are high.  $\text{SO}_2$  can also form sulfates in the atmosphere, a component of fine particulate matter.

The Puget Sound area has experienced a significant decrease in  $\text{SO}_2$  from sources such as pulp mills, cement plants and smelters in the last two decades.

In 1971, the EPA set an annual  $\text{SO}_2$  standard of 0.03 ppm and a 24-hour standard of 0.14 ppm that could not be exceeded more than once a year. EPA changed the  $\text{SO}_2$  standard in June of 2010 to a shorter-term (1-hour) standard of 0.075 ppm (75 ppb) and revoked the former annual and daily average standards. Historic comparisons to federal and Washington State standards can be seen in our 2009 data summary which is available upon request.

The 2010 standard is a 3-year average of the 99<sup>th</sup> percentile of the daily 1-hour maximum concentrations. Levels must be below 75 ppb. Sulfur dioxide levels at the Seattle Beacon Hill site have been below the 2010 standard from 2011–2022.


Figure 22 shows the maximum 3-year average of the 99<sup>th</sup> percentile of 1-hour maximum concentrations at Beacon Hill which have stayed within the standard.

Additional information on  $\text{SO}_2$  is available at <https://www.epa.gov/so2-pollution>.

**Figure 22: Sulfur Dioxide (SO<sub>2</sub>) 1-Hour Maximum Concentrations (3-Year Average of the 99th Percentile) for the Puget Sound Region**

### Sulfur Dioxide Estimated Design Values

(3-year average of 99th percentile of 1-Hour average daily maximum concentrations)



Note: 2011 was the first year that the Design Value has been calculated and compared to the revised primary SO<sub>2</sub> standard.

\* Indicates an estimate based on incomplete data. Data less than 75% for a quarter at Beacon Hill in 2016 & 2017.

### Lead

Lead is a highly toxic metal that was used for many years in household products such as paints, transportation fuel, and industrial chemicals. Now that lead has been banned from paint and most fuels, the greatest sources of lead emissions, nationally, are industrial processes (particularly primary and secondary lead smelters) and battery manufacturers. And while lead has been removed from fuel for large aircraft, lead found in aviation gasoline (avgas), used by small aircraft, remains a concern nationally.

People and animals are mainly exposed to lead by breathing it in and ingesting it in food, water, soil, or dust. Lead accumulates in the blood, bones, muscles, and fat. Infants and young children are especially sensitive to even low levels of lead. Lead can have health effects ranging from behavioral problems and learning disabilities to seizures and death.

Since the phase-out of lead in most fuels and the closure of the Harbor Island secondary lead smelter in Seattle in 1984, levels of lead in ambient air have decreased substantially. For a historic look at the Puget Sound region's lead levels, please see page 87 of the 2007 Air Quality Data Summary which is available [here](#).

In October 2008, EPA strengthened the lead standard from  $1.5 \mu\text{g}/\text{m}^3$  to  $0.15 \mu\text{g}/\text{m}^3$  (rolling three-month average).<sup>16</sup> As part of this rulemaking, EPA initiated a pilot lead monitoring program that focuses on lead from aviation gasoline at small airports, including two in our region. Results are available here: <https://fortress.wa.gov/ecy/publications/SummaryPages/1302040.html>. EPA maintained this level in its 2016 review of the lead standard.

For additional information on lead, visit <https://www.epa.gov/lead-air-pollution>.

---

<sup>16</sup>US EPA, National Ambient Air Quality Standard for Lead, Final Rule. Federal Register, November 12, 2008; <http://www.gpo.gov/fdsys/pkg/FR-2008-11-12/pdf/E8-25654.pdf>

### Visibility


Visibility data is presented as an indicator of air quality. Visibility is explained in terms of visual range and light extinction. *Visual range* is the maximum distance, usually in miles or kilometers, at which a black object is visible against the horizon. *Light extinction* is the sum of light scattering and light absorption by fine particles and gases in the atmosphere. The more light extinction, the shorter the visual range.

Reduced visibility is caused by weather such as clouds, fog, rain, and air pollution, including fine particles and gases. The major contributor to reduced visual range is fine particulate matter ( $PM_{2.5}$ ), which is present near the ground and can be transported aloft and may remain suspended for a week or longer. Figures 23 and 24 show visibility for the overall Puget Sound area, as well as 12-month moving average for King, Kitsap, Pierce and Snohomish Counties. Visibility on these graphs, in units of miles, is determined by continuous nephelometer monitoring. The nephelometer measures light scattering due to particulate matter ( $b_{sp}$ ), and this value is converted into estimates of visibility in miles. Nephelometer data are shown on page A-13 of the Appendix.


The red line represents the monthly average visibility. The large fluctuations are due to seasonal variability. The blue line shows the average of the previous 12-months. This moving average reduces seasonal variation and allows longer-term trends to be observed. The moving average shows that the visibility for the Puget Sound area has steadily increased (improved) over the last decade with some year-to-year variability. For the 24-year period from December 1990 through December 2022, the 12-month moving average increased from 48 miles to 86 miles.

For additional information on visibility, visit <https://www.epa.gov/visibility>.

**Figure 23: Puget Sound Visibility**



**Figure 24: County-wide Visibility**



### Air Toxics

“Air toxics” are air pollutants known or suspected to cause health problems. Potential health effects include cancer, birth defects, lung damage, immune system damage, and nerve damage.<sup>17,18</sup> The Agency considers over 400 different air pollutants air toxics.

This section presents a relative ranking of these toxics based on potential cancer risks, as well as trends over time. We provide a short description of each air toxic of concern, including their health effects and sources.

In 2021-2022, we completed an in-depth study of six locations in Tacoma and Seattle and finished a report in December 2023. At the time of this data summary, we did not have a reference yet to the study, but we will include it in subsequent data summaries.

Ecology monitors for air toxics at the Seattle Beacon Hill site. The Beacon Hill site is one of 27 EPA-sponsored National Air Toxic Trends Sites across the country.<sup>19</sup> As in previous years, Ecology monitored toxics every six days. The 2006 dataset is incomplete due to relocation of the Beacon Hill site that year; and the 2020 dataset is the average of only seven months and does not include some summer samples as the air toxics sampling could not be conducted from mid-March to July due to COVID-19. For general information on air toxics, see [www.pscleanair.gov/162/Air-Toxics](http://www.pscleanair.gov/162/Air-Toxics). Air toxics statistical summaries are provided starting on page A-17 of the Appendix.

### Relative ranking based on cancer risk & unit risk factors

Table 5 below ranks 2022 air toxics from the Seattle Beacon Hill monitoring site according to mean potential cancer risk per million people. It shows monitored pollutants ranked from highest concern (#1) to lowest, based on ambient concentrations multiplied by unit risk factors. A unit risk factor takes into account how toxic or carcinogenic a pollutant is. Cancer risk estimates are shown here to provide a meaningful basis of comparison between pollutants and are not intended to represent any one community’s or individual’s exposure.

Since the release of the 2018 Air Quality Data Summary, Ecology released an updated list of cancer risk factors, known as the Ambient Source Impact Level (ASIL) table.<sup>20</sup> The main reason for this update was to align the ASIL table with the latest scientific data. Ecology does not determine the unit risk factors that are the basis of the table, but rather uses values from three authoritative sources: the

<sup>17</sup>US EPA, Hazardous Air Pollutants: <https://www.epa.gov/haps>.

<sup>18</sup>US EPA, Risk Assessment for Toxic Air Pollutants: A Citizen’s Guide: [https://www3.epa.gov/airtoxics/3\\_90\\_024.html](https://www3.epa.gov/airtoxics/3_90_024.html).

<sup>19</sup> <https://www3.epa.gov/ttnamti1/natts.html>

<sup>20</sup> Washington Administrative Code Section 173-460-150; <https://apps.leg.wa.gov/wAc/default.aspx?cite=173-460-150>

## 2022 Air Quality Data Summary

EPA's Integrated Risk Information System (IRIS); the California Office of Environmental Health Hazard Assessment (OEHHA), and the U.S. Agency for Toxic Substances and Disease Registry (ATSDR). Many unit risk factors were updated by these agencies between the last ASIL update, in 2009, and 2019, leading to sometimes significant changes in the predicted risk. In some cases, the predicted risks have increased, such as with ethylene oxide, while in others, they have decreased, such as with carbon tetrachloride. These changes do not mean that the inherent risk of these pollutants has changed, but rather that our scientific understanding of the risk has improved, and we now have better estimates of their carcinogenicity.

Potential cancer risk is an estimate of the number of potential additional cancers (out of a population of one million) that may develop from exposure to air toxics over a lifetime (set at 70 years). A risk threshold of one in one million is commonly used as a screening value and is used here.<sup>21</sup> For details on how air toxics were ranked, please see page A-18 in the Appendix.

Risks presented in this table are based on annual average ambient (outside) concentrations. Risks based on 95<sup>th</sup> percentile concentrations (a more conservative statistic than presented in

Table 5) are presented on page A-19 of the Appendix. Page A-19 also lists the frequency (percentage) of samples that were over the cancer screening level of one in a million risk.

Two of the air toxics that present the greatest potential health risk in the Puget Sound area, diesel particulate matter and wood smoke particulate, are not included in the table. No direct monitoring method currently exists for these toxics. Modeling for these air toxics was not conducted for this report, however, the Agency has estimated the cancer risk for these parameters in recent studies.<sup>22,23</sup> Diesel Exhaust risk estimates can range from 400–600 per million in near-road and industrial areas; whereas the cancer risk from wood smoke can range from 20–30 per million in wood smoke impacted areas.

---

<sup>21</sup>US EPA, A Preliminary Risk-Based Screening Approach for Air Toxics Monitoring Datasets. EPA-904-B-06-001, Version 2, October 2010; [https://www.epa.gov/sites/production/files/2020-01/documents/air\\_1\\_preliminary\\_risk-based\\_screening\\_approach\\_p1009a7c.pdf](https://www.epa.gov/sites/production/files/2020-01/documents/air_1_preliminary_risk-based_screening_approach_p1009a7c.pdf)

<sup>22</sup> Puget Sound Clean Air Agency. 2010. Tacoma and Seattle Area Air Toxics Evaluation. <https://www.pscleanair.gov/DocumentCenter/View/2361>

<sup>23</sup> Puget Sound Clean Air Agency. 2018. Near-road air toxics study in the Chinatown-International District. <https://www.pscleanair.gov/DocumentCenter/View/3398/Air-Toxics-Study-in-the-Chinatown-International-District-Full-Report>

**Table 5: 2022 Beacon Hill Air Toxics Ranking**

(Average Potential Cancer Risk Estimate per 1,000,000)

Air Toxic	Rank	Average Potential Cancer Risk <sup>a</sup>
Ethylene oxide	1	926 <sup>b</sup>
Formaldehyde	2	7
Hexavalent Chromium	3	5 <sup>c</sup>
Benzene	3	4
Carbon tetrachloride	4	3
Acetaldehyde	5	3
Chloroform	6	2
Arsenic (PM <sub>10</sub> )	7	2
Acrolein	8	1
1,3-Butadiene	9	1
Naphthalene	10	1
Acrylonitrile	11	<1
Cadmium (PM <sub>10</sub> )	12	<1

<sup>a</sup> Risk based on unit risk factors as adopted in Washington State Acceptable Source Impact Level Table, 2019 update (WAC 173-460-150)<sup>24</sup>

<sup>b</sup> Ethylene oxide levels are very near detection limits. These values may not represent the risk accurately and may be subject to method bias.

<sup>c</sup> Sampling for hexavalent chromium was discontinued in 2013 and the included estimate is based on 2013 data using 2019 ASILs.

PM<sub>10</sub> = fine particles less than 10 micrometers in diameter

### Health effects other than cancer

Air toxics can also have chronic non-cancer health effects. These include respiratory, cardiac, immunological, nervous system, and reproductive system effects.

To determine non-cancer health risks, we compared each air toxic to its chronic reference exposure level, as established by California EPA (the most comprehensive dataset available)<sup>25</sup>. A chronic

<sup>24</sup>Washington State Administrative Code WAC 173-460-150, [apps.leg.wa.gov/WAC/default.aspx?cite=173-460-150](https://apps.leg.wa.gov/WAC/default.aspx?cite=173-460-150)

<sup>25</sup> <https://oehha.ca.gov/air/general-info/oehha-acute-8-hour-and-chronic-reference-exposure-level-rel-summary>

## 2022 Air Quality Data Summary

reference exposure level (chREL) is considered a safe level of continuous exposure to an individual air toxic for non-cancer health effects.

Only one air toxic, acrolein, failed the screen for non-cancer chronic health effects, with measured concentrations consistently exceeding the chREL. Acrolein irritates the lungs, eyes, and nose, and is a combustion by-product.<sup>26</sup> A table of reference concentrations and hazard indices for each air toxic measured in the last year with a hazard index greater than zero is on page A-20 of the Appendix. A hazard index is the concentration of a pollutant (either mean or other statistic) divided by the reference concentration. Typically, no adverse non-cancer health effects for that pollutant are associated with a hazard index less than 1, although it is important to consider that people are exposed to many pollutants at the same time. We did not explore acute non-cancer health effects, which are based on 1-hour measurements, because the Beacon Hill air toxics measurements are made on 24-hour samples.

### Air toxics trends

Trends in annual average cancer risks are shown on the following pages for the highest-ranked air toxics measured at Seattle Beacon Hill from 2000 to 2022. For many air toxics, our analysis of the trends shows a statistically significant decrease in annual average concentrations. EPA has not set ambient air standards for air toxics, so graphs do not include reference lines for federal standards. A statistical summary of the trends shown on the following pages can be found on page A-21 of the Appendix.

---


<sup>26</sup>EPA, Acrolein Hazard Summary: <https://www.epa.gov/sites/production/files/2016-08/documents/acrolein.pdf>.

### Ethylene Oxide

The EPA lists ethylene oxide as a known human carcinogen. Ethylene oxide inhalation is associated with increased risk of blood cancers and of breast cancer in women.<sup>27</sup> Its main use is as a chemical intermediate in the production of ethylene glycol (antifreeze), but it is also used as a fumigating agent for spices and cosmetics, and a sterilizing agent for medical supplies. Ethylene oxide's 2022 average potential cancer risk estimate at Seattle Beacon Hill was 926 in one million.

It was added to the suite of air toxics measured at Beacon Hill mid-year in 2018. In the 2018 ASIL update, ethylene oxide's unit risk factor was made more stringent by a factor of 57 (from 0.0114 to 0.0002). In 2019, ethylene oxide sampling across the country showed that the Seattle Beacon Hill monitor was one of the lowest across the country.<sup>28</sup> Although the value went up in 2022, it is still within the uncertainty of the measurements. As the ASIL is well below the detection limit, the annual average value has a significant degree of uncertainty (well over 100 per million potential cancer risk). The Agency is working alongside the EPA to determine key sources of ethylene oxide and reduction strategies.<sup>29</sup>

**Figure 25: Ethylene Oxide Annual Average Potential Cancer Risk at Beacon Hill, 2019–2022**



<sup>27</sup> EPA Hazard Summary: <https://www.epa.gov/sites/production/files/2016-09/documents/ethylene-oxide.pdf>

<sup>28</sup> EPA 2019. Map of NATTS/UTAMP Sites. [https://www.epa.gov/sites/production/files/2019-11/documents/map\\_of\\_natts\\_uatmp.pdf](https://www.epa.gov/sites/production/files/2019-11/documents/map_of_natts_uatmp.pdf)

<sup>29</sup> EPA 2020. EPA's Work to Understand Background Levels of Ethylene Oxide. <https://www.epa.gov/system/files/documents/2021-10/background-eto-explainer-document.pdf>


### Formaldehyde

The EPA lists formaldehyde as a probable human carcinogen. Formaldehyde inhalation is also associated with eye, nose, throat, and lung irritation.<sup>30</sup> Sources of ambient formaldehyde include automobiles, trucks, wood burning and other combustion. Formaldehyde's 2022 average potential cancer risk estimate at Beacon Hill was 7.5 in one million.

The sharp increase in average formaldehyde concentration in 2003 was due to nine anomalous sampling days in July 2003 when levels were roughly ten times the normal levels. It is possible that a local formaldehyde source was present at the Beacon Hill reservoir during this month and inadvertently affected the monitors.

Agency efforts that target vehicle exhaust and wood stove emission reductions also reduce formaldehyde emissions. Since 2000, we have found a statistically significant drop in risk from formaldehyde at an average rate of about 0.31 per million per year; however, the risk had been increasing in the recent years.

**Figure 26: Formaldehyde Annual Average Potential Cancer Risk at Beacon Hill, 2000–2022**



<sup>30</sup>EPA Hazard Summary; <https://www.epa.gov/sites/production/files/2016-09/documents/formaldehyde.pdf>.

### Hexavalent Chromium


Chromium is present in two chemical states (trivalent and hexavalent) in our air. Trivalent chromium occurs naturally, while hexavalent comes from human activities and is much more toxic. EPA lists hexavalent chromium as a known carcinogen, associated primarily with lung cancer. Hexavalent chromium is often abbreviated as chromium +6 or chromium (VI).

Exposure to hexavalent chromium is also associated with adverse respiratory, liver, and kidney effects.<sup>31</sup> Sources of hexavalent chromium include industrial processes such as chrome electroplating, as well as combustion of distillate oil, green glass production, and combustion of gasoline and diesel fuels (car, truck, and bus exhaust).

Due to the significant cost of monitoring for this pollutant, monitoring for total suspended particulate (TSP) hexavalent chromium was stopped in June 2013. The 2013 estimated average potential cancer risk for hexavalent chromium at Beacon Hill was 3 in one million based on the first half of the year.

In some years, up to 20% of the samples were below method detection limits. For the trend below, we used Kaplan-Meier analysis to estimate the annual means, as this method is designed to overcome bias from samples below the detection limit and other forms of censored data. Since 2000, we found a statistically significant drop in risk from hexavalent chromium at an average rate of about 0.7 per million per year. The Agency's permitting program works with and regulates industrial chromium plating operations to reduce hexavalent chromium emissions.

**Figure 27: Hexavalent Chromium Annual Average Potential Cancer Risk at Beacon Hill, 2005–2013**




<sup>31</sup>EPA Hazard Summary: <https://www.epa.gov/sites/production/files/2016-09/documents/chromium-compounds.pdf>.

### Benzene

The EPA lists benzene as a known human carcinogen. Benzene inhalation is also linked with blood, immune and nervous system disorders.<sup>32</sup> This air toxic comes from a variety of sources, including car/truck exhaust, wood burning, evaporation of industrial solvent and other combustion. Benzene's 2022 average potential cancer risk estimate at Beacon Hill was 4 in one million.

Benzene levels are likely decreasing in our area due to factors including less automobile pollution with cleaner vehicles coming into the fleet, better fuels, and fewer gas station emissions due to better compliance (vapor recovery at the pump and during filling of gas station tanks). We have found a statistically significant drop in risk from benzene at an average rate of about 0.34 per million per year since 2000.

**Figure 28: Benzene Annual Average Potential Cancer Risk at Beacon Hill, 2000–2022**




<sup>32</sup>EPA Hazard Summary; <https://www.epa.gov/sites/production/files/2016-09/documents/benzene.pdf>.

### Arsenic

EPA lists arsenic as a known carcinogen. Exposure to arsenic is also associated with skin irritation and liver and kidney damage.<sup>33</sup> Arsenic is used to treat wood and in colored glass. Combustion of distillate oil is also a source of arsenic in the Puget Sound area. Arsenic's 2022 average potential cancer risk estimate at Beacon Hill was 2 in one million. Since 2000, we found a statistically significant drop in risk from arsenic at an average rate of about 0.05 per million per year.

The Agency's permitting program works with and regulates industrial users of arsenic to reduce emissions. Illegal burning, especially of treated wood, can also contribute to arsenic emissions in our area.

**Figure 29: Arsenic Annual Average Potential Cancer Risk at Beacon Hill, 2003–2022**




<sup>33</sup>EPA Hazard Summary; <https://www.epa.gov/sites/production/files/2016-09/documents/arsenic-compounds.pdf>.

### Acetaldehyde

The EPA lists acetaldehyde as a probable human carcinogen. Acetaldehyde inhalation is also associated with irritation of eyes, throat and lungs, and long-term effects similar to those of alcoholism.<sup>34</sup> Main sources of acetaldehyde include wood burning and car/truck exhaust. Acetaldehyde's 2022 average potential cancer risk estimate at Beacon Hill was 3 in one million.

Agency efforts that target vehicle exhaust and wood stove emission reductions also reduce acetaldehyde emissions. Since 2000, we have found a statistically significant drop in risk from acetaldehyde at an average rate of about 0.1 per million per year. However, recent years have shown a slight increase.

**Figure 30: Acetaldehyde Annual Average Potential Cancer Risk at Beacon Hill, 2000–2022**




<sup>34</sup>EPA Hazard Summary; <https://www.epa.gov/sites/production/files/2016-09/documents/acetaldehyde.pdf>.

### Carbon Tetrachloride

The EPA lists carbon tetrachloride as a probable human carcinogen. Carbon tetrachloride inhalation is also associated with liver and kidney damage.<sup>35</sup> It was widely used as a solvent in both industry and consumer applications and was banned from consumer use in 1995. Trace amounts are still emitted by wastewater treatment plants. Carbon tetrachloride is relatively ubiquitous, has a long half-life, and occurs in similar concentrations in urban and rural areas. Carbon tetrachloride's 2022 average potential cancer risk estimate at Beacon Hill was 3 in one million.

The Agency does not target efforts at reducing carbon tetrachloride emissions, as carbon tetrachloride has already been banned. We did not find a statistically significant trend in carbon tetrachloride levels over time.

**Figure 31: Carbon Tetrachloride Annual Average Potential Cancer Risk at Beacon Hill, 2000–2022**




<sup>35</sup>EPA Hazard Summary; <https://www.epa.gov/sites/production/files/2016-09/documents/carbon-tetrachloride.pdf>.

### Chloroform

The EPA lists chloroform as a probable human carcinogen. Chloroform inhalation is associated with central nervous system effects and liver damage.<sup>36</sup> Main sources of chloroform are water treatment plants and reservoirs.<sup>37</sup> Because the Beacon Hill monitoring site is located at the Beacon Hill reservoir, which was uncovered prior to 2009, the chloroform measurements from 2000 through 2008 may be higher than expected for most of our region. However, the reservoir underwent a major renovation in 2008-09 and is now completely enclosed, possibly at least partially explaining the drop in chloroform levels around that time. Chloroform's 2022 average potential cancer risk estimate at Beacon Hill was 2 in one million.

The Agency does not prioritize efforts to reduce chloroform emissions, as it does not likely present risk in areas other than those directly adjacent to reservoirs, the majority of which have been covered in accordance with a 2006 federal regulation on drinking water protection.<sup>38</sup> Since 2000, we have found a statistically significant drop in risk from chloroform at an average rate of about 0.17 per million per year.

**Figure 32: Chloroform Annual Average Potential Cancer Risk at Beacon Hill, 2000–2022**



<sup>36</sup>EPA Hazard Summary; <https://www.epa.gov/sites/production/files/2016-09/documents/chloroform.pdf>.

<sup>37</sup>Seattle Public Utilities. 2018 Water Quality Analysis shows detectable levels of trihalomethanes in treated drinking water, which is stored in reservoirs (trihalomethanes include chloroform, dichlorobromomethane, dibromochloromethane, and bromoform); [https://www.seattle.gov/Documents/Departments/SPU/Services/Water/Water\\_Quality\\_Report\\_2018.pdf](https://www.seattle.gov/Documents/Departments/SPU/Services/Water/Water_Quality_Report_2018.pdf).

<sup>38</sup>Long Term 2 Enhanced Surface Water Treatment Rule; <https://www.epa.gov/dwreginfo/long-term-2-enhanced-surface-water-treatment-rule-documents>


### Ethylene Dichloride

EPA lists ethylene dichloride as a probable human carcinogen. It is primarily used as a solvent in the production of other chemicals like vinyl chloride. It is also added to leaded gasoline, but this is expected to be a very minor source, as leaded gas for on-road vehicle use was phased out in 1996.<sup>39,40</sup> We estimated ethylene dichloride's 2022 average potential cancer risk estimate at Beacon Hill at 2 in one million.

There is no useful trend information for this air toxic since most of the measurements are near the practical quantitation limit of the analytical method. That is, 44% of the samples in 2022 were within twice the method detection limit. Additionally, in prior years, most of the samples were also below the quantitation limits. In the years for which we have ethylene dichloride data, the detection limit for this air toxic is typically near the one in a million potential cancer risk level.

The Agency's permitting program works with and regulates industrial producers of ethylene dichloride to reduce emissions.

**Figure 33: Ethylene Dichloride Annual Average Potential Cancer Risk at Beacon Hill, 2014–2022**



<sup>39</sup> EPA Hazard Summary, <https://www.epa.gov/sites/production/files/2016-09/documents/ethylene-dichloride.pdf>.


<sup>40</sup> US Energy Information Administration: Gasoline and the Environment; [https://www.eia.gov/energyexplained/index.php?page=gasoline\\_environment](https://www.eia.gov/energyexplained/index.php?page=gasoline_environment)

### 1,3-Butadiene

The EPA lists 1,3-butadiene as a known human carcinogen. 1,3-butadiene inhalation is also associated with neurological effects.<sup>41</sup> Primary sources of 1,3-butadiene include cars, trucks, buses, and wood burning. 1,3-butadiene's 2022 average potential cancer risk estimate at Beacon Hill was 1 in one million.

Agency efforts that target vehicle exhaust and wood stove emission reductions also reduce 1,3-butadiene emissions. Since 2000, we have found a statistically significant drop in risk from 1,3-butadiene at an average rate of about 0.1 per million per year.

**Figure 34: 1,3-Butadiene Annual Average Potential Cancer Risk at Beacon Hill, 2000–2022**




<sup>41</sup>EPA Hazard Summary: <https://www.epa.gov/sites/production/files/2016-08/documents/13-butadiene.pdf>.

### Naphthalene

EPA lists naphthalene as a possible human carcinogen. Naphthalene is also associated with respiratory effects and retina damage.<sup>42</sup> Local sources of naphthalene include combustion of wood and heavy fuels. Naphthalene's 2022 average potential cancer risk estimate at Beacon Hill was at 1 in one million.

The Agency works with and regulates wood burning through burn bans and wood stove replacement programs to reduce naphthalene emissions. Since 2014, we have found a statistically significant drop in risk from naphthalene at an average rate of about 0.08 per million per year. Monitoring for naphthalene and other polycyclic aromatic hydrocarbons started in 2008.

**Figure 35: Naphthalene Annual Average Potential Cancer Risk at Beacon Hill, 2008–2022**




<sup>42</sup>EPA Hazard Summary; <https://www.epa.gov/sites/production/files/2016-09/documents/naphthalene.pdf>.

### **Tetrachloroethylene**

EPA lists tetrachloroethylene as a probable human carcinogen. It is widely used for dry-cleaning fabrics and textile processing. It is also used for vapor degreasing in metal cleaning process. Chronic exposure to tetrachloroethylene may affect kidney, liver, immune system, and neurological behavior. We estimated tetrachloroethylene's 2022 average potential cancer risk estimate at Beacon Hill was less than one in one million.

Since 2007, we have found a statistically significant drop in risk from tetrachloroethylene at an average rate of about 0.04 per million per year. The Agency's compliance program works with and regulates dry cleaners and other producers of tetrachloroethylene to reduce emissions.

**Figure 36: Tetrachloroethylene Annual Average Potential Cancer Risk at Beacon Hill, 2000–2022**




### Ethylbenzene

EPA lists ethylbenzene as a Group D pollutant, which is not classifiable as to human carcinogenicity due to limited information available.<sup>43</sup> Chronic exposure to ethylbenzene may affect the blood, liver, and kidneys. Local sources of ethylbenzene are from fuels, asphalt, and naphtha. It is also used in styrene production. Ethylbenzene's 2022 average potential cancer risk estimate at Beacon Hill was less than one in one million.

We did not find a statistically significant trend in ethylbenzene levels over the 2007–2022 timeframe for which we have data. The Agency works with and regulates solvent-using businesses to reduce ethylbenzene emissions.

**Figure 37: Ethylbenzene Annual Average Potential Cancer Risk at Beacon Hill, 2007–2022**



<sup>43</sup>EPA Hazard Summary: <https://www.epa.gov/sites/production/files/2016-09/documents/ethylbenzene.pdf>.


### Cadmium

EPA lists cadmium as a probable human carcinogen. Cadmium exposures are also associated with kidney damage.<sup>44</sup> Combustion of distillate oil is a main source of cadmium in the Puget Sound area.

Cadmium's 2022 average potential cancer risk estimate at Beacon Hill was less than 1 in one million. Our trend is affected by a number of factors, including the fact that over half the samples in 2010 were below the detection limits and thus we did not have sufficient data to make a comparable average. Extremely high outlier results on 11/18/13 and 9/8/14 resulted in high average concentrations in each of those respective years. On those days, no other metal concentrations were statistical outliers compared to their respective annual variability. With the outliers excluded for 2013 and 2014, the estimated annual potential cancer risks for those years would be < 1. With or without the outliers included, we found no statistically significant trend for cadmium.

The Agency's permitting program works with and regulates industrial producers of cadmium to reduce emissions.

**Figure 38: Cadmium Annual Average Potential Cancer Risk at Beacon Hill, 2003–2022**



<sup>44</sup>EPA Hazard Summary; <https://www.epa.gov/sites/production/files/2016-09/documents/cadmium-compounds.pdf>.

## Definitions

### General Definitions

#### Air Quality Index

**Table 6: 2022 Calculation and Breakpoints for the Air Quality Index (AQI)**

Breakpoints for Criteria Pollutants							AQI Categories	
O <sub>3</sub> (ppm) 8-hour	O <sub>3</sub> (ppm) 1-hour <sup>(a)</sup>	PM <sub>2.5</sub> ( $\mu\text{g}/\text{m}^3$ ) 24 hour	PM <sub>10</sub> ( $\mu\text{g}/\text{m}^3$ ) 24 hour	CO (ppm) 8 hour	SO <sub>2</sub> <sup>(c)</sup> (ppb) 1 hour	NO <sub>2</sub> (ppb) 1 hour	AQI value	Category
0.000–0.054	—	0.0–12.0	0–54	0.0–4.4	0–35	0–53	0–50	Good
0.055–0.070	—	12.1–35.4	55–154	4.5–9.4	36–75	54–100	51–100	Moderate
0.071–0.085	0.125–0.164	35.5–55.4	155–254	9.5–12.4	76–185	101–360	101–150	Unhealthy for sensitive groups
0.086–0.105	0.165–0.204	55.5–150.4	255–354	12.5–15.4	(186–304) <sup>(d)</sup>	361–649	151–200	Unhealthy
0.106–0.200	0.205–0.404	150.5–250.4	355–424	15.5–30.4	(305–604) <sup>(d)</sup>	650–1249	201–300	Very unhealthy
(b)	0.405–0.504	250.5–350.4	425–504	30.5–40.4	(604–804) <sup>(d)</sup>	1250–1649	301–400	Hazardous
(b)	0.505–0.604	350.5–500.4	505–604	40.5–50.4	(805–1004) <sup>(d)</sup>	1650–2049	401–500	

<sup>(a)</sup>Areas are generally required to report the AQI based on 8-hour ozone values. However, there are a small number of areas where an AQI based on 1-hour ozone values would be safer. In these cases, in addition to calculating the 8-hour ozone value, the 1-hour ozone value may be calculated, and the greater of the two values reported.

<sup>(b)</sup>8-hour O<sub>3</sub> values do not define higher AQI values (above 300). AQI values above 300 are calculated with 1-hour O<sub>3</sub> concentrations.

<sup>(c)</sup>EPA changed the SO<sub>2</sub> standard on June 22, 2010 to be based on an hourly maximum instead of a 24-hour and annual average.

<sup>(d)</sup>1-hour SO<sub>2</sub> values do not define higher AQI values ( $\geq 200$ ). AQI values of 200 or greater are calculated with 24-hour SO<sub>2</sub> concentrations.

For more information on the AQI, see [airnow.gov/index.cfm?action=aqibasics.aqi](http://airnow.gov/index.cfm?action=aqibasics.aqi).

#### Air shed

A geographic area that shares the same air, due to topography, meteorology and climate.

#### Air Toxics

Air toxics are broadly defined as over 400 pollutants that the Agency considers potentially harmful to human health and the environment. These pollutants are listed in the Washington Administrative Code at [apps.leg.wa.gov/WAC/default.aspx?cite=173-460-150](http://apps.leg.wa.gov/WAC/default.aspx?cite=173-460-150). Hazardous air pollutants (see below) are checked on this list to identify them as a subset of air toxics. Air toxics are also called Toxic Air Contaminants (TAC) under Agency Regulation III.

#### Criteria Air Pollutant (CAP)

The Clean Air Act of 1970 defined *criteria pollutants* and provided EPA the authority to establish ambient concentration standards for these criteria pollutants to protect public health. EPA periodically revises the original concentration limits and methods of measurement, most recently in 2011. The six criteria air pollutants are: particulate matter (10 micrometers and 2.5

micrometers), ozone, nitrogen dioxide, carbon monoxide, sulfur dioxide and lead. See Appendix page A-22 for more information.

### **ppm, ppb (parts per million, or parts per billion)**

A unit of concentration used for many air pollutants. A ppm (ppb) means one molecule of the pollutant per million (or billion) molecules of air.

### **Hazardous Air Pollutant (HAP)**

A *hazardous air pollutant* is an air contaminant listed in the Federal Clean Air Act, Section 112(b). EPA currently lists 187 pollutants as HAPs at <https://www.epa.gov/haps/initial-list-hazardous-air-pollutants-modifications>.

### **Temperature Inversions**

Air temperature usually decreases with altitude. On a sunny day, air near the surface is warmed and is free to rise. The warm surface air can rise to altitudes of 4,000 feet or more and is dispersed (or mixed) into higher altitudes. In contrast, on clear nights with little wind, the surface can cool rapidly (by 10 degrees or more), which also cools the air just above the surface. The air aloft does not cool, which creates a very stable situation where the warm air aloft effectively caps the cooler air below. This process limits mixing to just a few hundred feet or less. This situation is called a temperature inversion and allows for pollutants to accumulate to high concentrations.

### **Unit Risk Factor (URF)**

A unit risk factor is a measure of a pollutant's cancer risk based on a 70-year inhalation exposure period. The units are risk/concentration. Unit risk factors are multiplied by concentrations to estimate potential cancer risk.

### **Visibility/Regional Haze**

Visibility is often explained in terms of visual range and light extinction. *Visual range* is the maximum distance (usually miles or kilometers) a black object can be seen against the horizon. *Light extinction* is the sum of light scattering and light absorption by fine particles and gases in the atmosphere. The more light extinction, the shorter the visual range. Reduced visibility (or visual range) is caused by weather (clouds, fog, and rain) and air pollution (fine particles and gases).

### **Volatile Organic Compound (voc)**

An organic compound that participates in atmospheric photochemical reactions. This excludes compounds determined by EPA to have negligible photochemical reactivity.



PUGET SOUND

**Clean Air Agency**

1904 Third Avenue, Suite 105  
Seattle, Washington 98101  
[www.pscleanair.gov](http://www.pscleanair.gov)

---

**2022**  
**Air Quality**  
**Data Summary Appendix**

December 2023

## Table of Contents

Monitoring Methods Used from 1999 to 2022 in the Puget Sound Air Shed.....	3
Historical Air Quality Monitoring Network.....	4
Burn Bans 1988 – 2022.....	9
Particulate Matter (PM <sub>2.5</sub> ) – Federal Reference Method.....	11
Particulate Matter (PM <sub>2.5</sub> ) – Federal Equivalent Methods.....	12
Particulate Matter (PM <sub>2.5</sub> ) – Continuous – Nephelometer.....	13
PM <sub>2.5</sub> Speciation Analytes Monitored in 2022 .....	14
PM <sub>2.5</sub> Black Carbon.....	15
Ozone (8-hour concentration) .....	16
2022 Beacon Hill Air Toxics Statistical Summary for Air Toxics ( <i>units in parts per billion</i> ) .....	17
2022 Beacon Hill Air Toxics Statistical Summary for Air Toxics ( <i>units in nanograms per cubic meter</i> ) .....	17
Estimates of Air Toxics Risk.....	18
2022 Air Toxics Unit Risk Factors .....	18
2022 Beacon Hill Potential Cancer Risk Estimates per 1,000,000 – 95 <sup>th</sup> Percentile.....	19
2022 Non-cancer Reference Concentrations (RfC) and Hazard Indices >1 .....	20
2000–2022 Air Toxics Trends Statistical Summary .....	21
Air Quality Standards and Health Goals.....	22

## Monitoring Methods Used from 1999 to 2022 in the Puget Sound Air Shed

<b>Pollutant Code</b>	<b>Measurement</b>	<b>Method</b>	<b>Units</b>
Bap	Light Absorption by Particles	Light Absorption by Aethalometer	bap (x 10 exp-4)/m
Bsp	Light Scattering by Particles	Nephelometer - Heated Inlet	bsp (x 10 exp-4)/m
CO	Carbon Monoxide	Gas Nondispersive Infrared Radiation	parts per million
NO <sub>x</sub>	Nitrogen Oxides (NO <sub>x</sub> )	Chemiluminescence	parts per million
	Nitric Oxide (NO)	Chemiluminescence	parts per million
	Nitrogen Dioxide (NO <sub>2</sub> )	Chemiluminescence	parts per million
NO <sub>Y</sub>	Reactive Nitrogen Compounds (NO <sub>x</sub> + other reactive compounds)	Chemiluminescence	parts per billion
O <sub>3</sub>	Ozone	UV Absorption	parts per million
Pb	Lead	Standard High Volume	micrograms per standard cubic meter
PM <sub>10</sub> ref	PM <sub>10</sub> Reference	Reference - Hi Vol Andersen/GMW 1200	micrograms per cubic meter
PM <sub>10</sub> bam	PM <sub>10</sub> Beta Attenuation	Andersen FH62I-N	micrograms per cubic meter
PM <sub>10</sub> teom	PM <sub>10</sub> Teom	R&P Mass Transducer	micrograms per cubic meter
PM <sub>2.5</sub> ref	PM <sub>2.5</sub> Reference	Reference—R&P Partisol 2025	micrograms per cubic meter
PM <sub>2.5</sub> bam	PM <sub>2.5</sub> Beta Attenuation	Andersen FH62I-N	micrograms per cubic meter
PM <sub>2.5</sub> teom	PM <sub>2.5</sub> Teom	R&P Mass Transducer	micrograms per cubic meter
PM <sub>2.5</sub> ls	PM <sub>2.5</sub> Nephelometer	Radiance Research M903 Nephelometer	micrograms per cubic meter
PM <sub>2.5</sub> bc	PM <sub>2.5</sub> Black Carbon	Light Absorption by Aethalometer	micrograms per cubic meter
RH	Relative Humidity	Continuous Instrument Output	percent
SO <sub>2</sub>	Sulfur Dioxide	UV Fluorescence	parts per million
Temp	Temperature	Continuous Instrument Output	degrees F
TSP	PM Total Hi-Vol	Standard High Volume	micrograms per standard cubic meter
Vsby	Visual Range	Light Scattering by Nephelometer	miles
Wind	Wind Speed/ Wind Direction	RM Young 05305 Wind Monitor AQ (old method)	miles per hour/degrees
	Wind Speed/ Wind Direction	Ultrasonic (new method)	miles per hour/degrees

## Historical Air Quality Monitoring Network

Station ID	Location	PM <sub>10</sub> Ref	PM <sub>10</sub> bam	PM <sub>10</sub> teom	PM <sub>2.5</sub> ref	PM <sub>2.5</sub> bam	PM <sub>2.5</sub> teom	PM <sub>2.5</sub> ls	PM <sub>2.5</sub> bc	O <sub>3</sub>	SO <sub>2</sub>	NO <sub>x</sub>	CO	b <sub>sp</sub>	Wind	Temp	AT	Vsby	Location
AO◎	Northgate, 310 NE Northgate Way, Seattle (ended Mar 31, 2003)												X						b, d, f
AQ	Queen Anne Hill, 400 W Garfield St, Seattle (photo/visibility included) (ended 3/18/2015)							X						X	X	X		X	a, d, f
AR◎	4th Ave & Pike St, 1424 4 <sup>th</sup> Ave, Seattle (ended Jun 30, 2006)												X						a, d
AS◎	5th Ave & James St, Seattle (ended Feb 28, 2001)												X						a, d
AU◎	622 Bellevue Way NE, Bellevue (ended Jul 30, 1999)												X						a, d
AZ	Olive Way & Boren Ave, 1624 Boren Ave, Seattle SPECIATION SITE (ended 8/6/2014)							X	X					X	X	X		X	a, d
BF◎	University District, 1307 NE 45th St, Seattle (ended Jun 30, 2006)												X						b, d
BK◎	10 <sup>th</sup> & Weller, Seattle SPECIATION SITE					●	X		●			●	●		●	●	●		a
BL	11675 44 <sup>th</sup> Ave S, Tukwila Allentown					●	X	●	●					●	●	●	●	●	b, e, f
BU◎	Highway 410, 2 miles E of Enumclaw (ended Sep 30, 2000)									X									c, e
BV	Sand Point, 7600 Sand Pt Way NE, Seattle (ended Aug 31, 2006)							X						X	X	X			b, d
BW◎/BZ◎	Beacon Hill, 15th S & Charlestown, Seattle SPECIATION SITE				●	●	X	X	X	●	●	●	●	X	●	●	●	X	b, d, f
CE	Duwamish, 4700/4752 E Marginal Way S, Seattle SPECIATION SITE	X		X	●	●	X	●	●		X			●	●	●	●	●	a, e
CG◎	Woodinville, 17401 133 <sup>rd</sup> Av NE, Woodinville (ended April 2010)							X						X					b, d, f
CW	James St & Central Ave, Kent	X		X	X	●	X	●	●					●	●	●		●	b, d
CX	17711 Ballinger Way NE, Lake Forest Park (ended Jun 4, 1999)	X	X											X	X			X	b, d, f
CZ	Aquatic Center, 601 143 <sup>rd</sup> Ave NE, Bellevue (ended May 31, 2006)						X	X						X				X	b, f

Station ID	Location	PM <sub>10</sub> Ref	PM <sub>10</sub> bam	PM <sub>10</sub> teom	PM <sub>2.5</sub> ref	PM <sub>2.5</sub> bam	PM <sub>2.5</sub> teom	PM <sub>2.5</sub> ls	PM <sub>2.5</sub> bc	O <sub>3</sub>	SO <sub>2</sub>	NO <sub>y</sub>	CO	b <sub>sp</sub>	Wind	Temp	AT	Vsby	Location
DA	South Park, 8025 10 <sup>th</sup> Ave S, Seattle (ended Dec 31, 2002)	X			X			X						X	X			X	b, e, f
DB	17171 Bothell Way NE, Lake Forest Park	X	X		X		X	●	X					●	X	X		●	b, d, f
DC●	305 Bellevue Way NE, Bellevue				X			●						●				●	a, d
DD	South Park, 8201 10 <sup>th</sup> Ave S, Seattle							●						●				●	b, e, f
DE●	City Hall, 15670 NE 85 <sup>th</sup> St, Redmond (ended Dec 14, 2005)				X			X						X				X	a, d
DF●	30525 SE Mud Mountain Road, Enumclaw				X			X		●				X	●	●		X	c
DG●	42404 SE North Bend Way, North Bend				X		X	●	●					●	X	●		●	c, d, f
DH●	2421 148 <sup>th</sup> Ave NE, Bellevue (ended 1/21/2010)													X					b, d
DK●	43407 212 <sup>th</sup> Ave SE, 2 mi west of Enumclaw (ended Sep 6, 2006)														X	X			c
DL●	NE 8th St & 108th Ave NE, Bellevue (ended March 4, 2003)													X					a, d
DN●	20050 SE 56 <sup>th</sup> , Lake Sammamish State Park, Issaquah									●					X	X			b, d
DP●	504 Bellevue Way NE, Bellevue (ended Sep 30, 1999)	X			X														a, d
DZ●	Georgetown, 6431 Corson Ave S, Seattle (ended August 31, 2002)											X	X		X				a, d, e, f
EA	Fire Station #12, 2316 E 11 <sup>th</sup> St, Tacoma (ended Dec 31, 2000)	X	X												X				a, e
EP	27th St NE & 54th Ave NE, Tacoma (ended Feb 29, 2000)	X										X			X				b, e, f
EQ	Tacoma Tideflats, 2301 Alexander Ave, Tacoma SPECIATION SITE	X	X	X	X	●	X	●	●		X			●	●	●	●	●	a, e
ER	South Hill, 9616 128 <sup>th</sup> St E, Puyallup (ended December 2, 2020)	X	X		X	X		X	X					X	X	X		X	b, f
ES	7802 South L St, Tacoma SPECIATION SITE				●	●	X	●	●					●	●	●	●	●	b, f
FF●	Tacoma Indian Hill, 5225 Tower Drive NE, northeast Tacoma (ended 12/10/20)													X	X				b, f

Station ID	Location	PM <sub>10</sub> Ref	PM <sub>10</sub> bam	PM <sub>10</sub> teom	PM <sub>2.5</sub> ref	PM <sub>2.5</sub> bam	PM <sub>2.5</sub> teom	PM <sub>2.5</sub> ls	PM <sub>2.5</sub> bc	O <sub>3</sub>	SO <sub>2</sub>	NO <sub>y</sub>	CO	b <sub>sp</sub>	Wind	Temp	AT	Vsby	Location
FG◎	Mt Rainier National Park, Jackson Visitor Center									●									c
FH◎	Charles L Pack Forest, La Grande (ended 9/30/2010)									X									c, f
FL◎	1101 Pacific Ave, Tacoma (ended Jun 30, 2006)												X						a, d
ID	Hoyt Ave & 26th St, Everett (ended Feb 29, 2000)										X				X				a, e, d
IG	Marysville JHS, 1605 7 <sup>th</sup> St, Marysville SPECIATION SITE	X	X		X	●	X	●	X					●	●	●	●	b, d	
IH	20935 59 <sup>th</sup> Place West, Lynnwood (ended Jun 8, 1999)	X		X										X	X		X	a, d	
II	6120 212 <sup>th</sup> St SW, Lynnwood (ended April 9, 2020)				X	X	X	X						X	X	X	X	b, d	
IK◎	14310 SE 12 <sup>th</sup> St, Bellevue						X	●						●				a, d	
JN◎	5810 196 <sup>th</sup> Street, Lynnwood (ended Jun 30, 2006)												X					a, d	
JO	Darrington High School, Darrington 1085 Fir St				X	●	X	●	X					●	●	●	●	d, f	
JP◎	2939 Broadway Ave, Everett (ended March 31, 2003)												X					a, d	
JQ◎	44th Ave W & 196 <sup>th</sup> St SW, Lynnwood (ended May 3, 2004)												X					a, d	
JS◎	Broadway & Hewitt Ave, Everett (ended May 21, 2000)												X					a, d	
PA◎	1802 S 36 <sup>th</sup> St, Tacoma					●	X		●			●			●	●	●	a, f	
QE	Meadowdale, 7252 Blackbird Dr NE, Bremerton (ended 5/1/2012)	X				X	X	X						X	X	X	X	b, f	
QF	Lions Park, 6th Ave NE & Fjord Dr, Poulsbo (ended Feb 29, 2000)													X				b, f	
QG	Fire Station #51, 10955 Silverdale Way, Silverdale (ended September 4, 2008)				X		X						X	X	X	X	a, d		
QK	Spruce, 3250 Spruce Ave, Bremerton					●	X	●					●	●	●	●	b, f		

Station ID	Location	PM <sub>10</sub> Ref	PM <sub>10</sub> bam	PM <sub>10</sub> teom	PM <sub>2.5</sub> ref	PM <sub>2.5</sub> bam	PM <sub>2.5</sub> teom	PM <sub>2.5</sub> ls	PM <sub>2.5</sub> bc	O <sub>3</sub>	SO <sub>2</sub>	NO <sub>y</sub>	CO	b <sub>sp</sub>	Wind	Temp	AT	Vsby	Location
RV@	Yelm N Pacific Road, 931 Northern Pacific Rd SE, Yelm									●									c, f
RZ	Gig Harbor, 9702 Crescent Valley Dr NW, Gig Harbor							X						X	X	X		X	f
TC	M St E, Auburn (ended April 7, 2020)							X						X	X	X		X	b
TR	Eatonville, 560 Center St, Eatonville							X						X	X	X		X	F
TS	1301 Yesler Way, Seattle								X								X		a, f
TT	602 S. Jackson St, Seattle								X								X		a, f
UB@	71 E Campus Dr, Belfair (ended Sep 30, 2004)									X									c
VK@	Fire Station, 709 Mill Road SE, Yelm (ended Oct 2005)									X									c, f
ZB	Auburn 29 <sup>th</sup> Street, 402 29 <sup>th</sup> Street, Auburn					●			●					●	●	●	●		b

---

◎	Station operated by Ecology	SO <sub>2</sub>	Sulfur Dioxide
RVO◎	Shading indicates station functioning	NO <sub>y</sub>	Nitrogen Oxides
●	Indicates parameter currently monitored	CO	Carbon Monoxide
X	Indicates parameter previously monitored	b <sub>sp</sub>	Light scattering by atmospheric particles (nephelometer)
PM <sub>10</sub> ref	Particulate matter <10 micrometers (reference)	Wind	Wind direction and speed
PM <sub>10</sub> bam	Particulate matter <10 micrometers (beta attenuation continuous)	Temp	Air temperature (relative humidity also measured at BW, IG, ES)
PM <sub>10</sub> teom	Particulate matter <10 micrometers (teom continuous)	AT	Air Toxics
PM <sub>2.5</sub> ref	Particulate matter <2.5 micrometers (reference)	VSBY	Visual range (light scattering by atmospheric particles)
PM <sub>2.5</sub> bam	Particulate matter <2.5 micrometers (beta attenuation continuous)	PHOTO	Visibility (camera)
PM <sub>2.5</sub> teom	Particulate matter <2.5 micrometers (teom-fdms continuous)	O <sub>3</sub>	Ozone (May through September)
PM <sub>2.5</sub> ls	Particulate matter <2.5 micrometers (light scattering nephelometer continuous)		
PM <sub>2.5</sub> bc	Particulate matter <2.5 micrometers black carbon (light absorption aethalometer)		
Location		e	Industrial
a	Urban Center	f	Residential
b	Suburban		
c	Rural		
d	Commercial		

## Burn Bans 1988 – 2022

1988	Jan 25 (0830) – Jan 28 (0830) Feb 5 (1630) – Feb 6 (0930) Dec 1 (1430) – Dec 2 (0800) Dec 4 (1430) – Dec 5 (1400) Dec 16 (1430) – Dec 18 (1430)	2004	None
1989	Jan 19 (1430) – Jan 20 (1430) Jan 24 (1430) – Jan 26 (0930) Feb 6 (1430) – Feb 8 (0930) Feb 10 (1430) – Feb 16 (0930) Nov 29 (1430) – Dec 2 (0930) Dec 22 (1430) – Dec 23 (1430)	2005	Feb 21 (1600) – Feb 28 (0800) Dec 9 (1700) – Dec 18 (1200)
1990	Jan 19 (1430) – Jan 21 (1430) Dec 7 (1430) – Dec 8 (0930) Dec 25 (1430) – Dec 27 (0815)* *(Dec 26 (1430) – Dec 27 (0815)) 2 <sup>nd</sup> Stage	2006	None
1991	Jan 5 (1430) – Jan 6 (0930) Jan 21 (1430) – Jan 24 (1500)* *(Jan 22 0930 – Jan 24 1500) 2 <sup>nd</sup> Stage Jan 29 (1430) – Jan 31 (0830) Dec 15 (1430) – Dec 17 (1430)* *(Dec 16 (1430) – Dec 17 (0930)) 2 <sup>nd</sup> Stage	2007	Jan 13 (1400) – Jan 16 (1500) Jan 28 (1400) – Jan 31 (1400) Dec 9 (1400) – Dec 11 (0930)
1992	Jan 8 (1430) – Jan 9 (0930) Jan 19 (1430) – Jan 20 (1430) Feb 5 (1000) – Feb 6 (1430) Nov 25 (1430) – Nov 26 (1430)	2008	Jan 23 (1400) – Jan 26 (1200)
1993	Jan 11 (1430) – Jan 13 (0830) Jan 15 (1430) – Jan 16 (0700) Jan 17 (1430) – Jan 19 (0600) Jan 31 (1430) – Feb 3 (0830) Dec 20 (1430) – Dec 21 (1430) Dec 26 (1430) – Dec 29 (0830)	2009	Jan 16 (1200) – Jan 24 (1200) Feb 3 (1400) – Feb 6 (0900) Dec 8 (1000) – Dec 13 (1000) Dec 23 (1600) – Dec 30 (1200)
1994	None	2010	Jan 28 (1200) – Jan 31 (1000) Dec 30 (1700) – 31 Dec (2400)* * continued to Jan 4 (1700)
1995	Jan 4 – Jan 7	2011	Jan 1 (0000) – Jan 4 (1700) Nov 30 (1700) – Dec 7 (1300) Dec 11 (1700) – Dec 14 (1600)
1996	Feb 14 (1430) – Feb 16 (1630)	2012	Jan 11 (1600) – Jan 14 (1000) Jan 27 (1200) – Jan 28 (1700) Feb 3 (1600) – Feb 6 (1600) Nov 25 (1300) – Nov 28 (0900) Dec 29 (1700) – Dec 31 (2400)* * continued to Jan 3 (1200)
1997	Nov 13 (1500) – Nov 15 (1500) Dec 4 (1500) – Dec 7 (1800)	2013	Jan 1 (0000) – Jan 3 (1200) Jan 12 (1300) – Jan 22 (1000) Nov 22 (1600) – Nov 29 (1000) Dec 7 (1400) – Dec 9 (1000) Dec 25 (1700) – Dec 26 (1100)
1998	None	2014	Jan 26 (1200) – Jan 27 (1000) Nov 14 (1700) – Nov 20 (0600) Nov 30 (1300) – Dec 2 (1200) Dec 30 (1600) – Dec 31 (2400)* * continued to Jan 3 (1200)
1999	Jan 5 (1400) – Jan 6 (1000) Dec 29 (1400) – Dec 31 (0600)	2015	Jan 1 (0000) – Jan 3 (1200) Jan 10 (1200) – Jan 10 (1900) Jan 11 (1200) – 12 Jan (1100) Nov 25 (1600) – Dec 1 (0800) 24 Dec (1600) – 25 Dec (0830)
2000	Feb 18 (1400) – Feb 20 (1000) Nov 15 (1700) – Nov 23 (0600)	2016	1 Jan (1300) – 4 Jan (0930) 7 Jan (1300) – 9 Jan (1200) 10 Jan (1300) – 11 Jan (0900) 15 Dec (1300) – 18 Dec (0900)
2001	Nov 8 (1400) – Nov 12 (1800)	2017	4 Jan (1800) – 7 Jan (1300) 11 Jan (1200) – 16 Jan (1700)
2002	Nov 1 (1500) – Nov 6 (0900) Nov 27 (1000) – Dec 4 (1000)		
2003	Jan 7 (1500) – Jan 9 (1300)		

	24 Jan (1400) – 25 Jan (1200) 2 Aug (1600) – 5 Aug (100) 8 Aug (1400) – 11 Aug (1400) 8 Dec (1400) – 13 Dec (1400) 22 Dec (1400) – 24 Dec (1200)	13 Jan (1300) – 16 Jan (1200)
	2020	None
	2021	None
2018	1 Jan (1400) – 2 Jan (1100) 20 Aug (1700) – 23 Aug (1300)	2022 13 Nov (1500) – 16 Nov (1300) 19 Nov (1300) – 22 Nov (0000) 16 Dec (1400) – 17 Dec (1300)
2019	1 Jan (1400) – 2 Jan (1000)	

## Particulate Matter (PM<sub>2.5</sub>) - Federal Reference Method

Micrograms per Cubic Meter

Reference Sampling Method: R&P Partisol 2025 Sampler – Teflon Filter

2022

Location	Number of Values	Quarterly Arithmetic Averages				Annual Mean	98 <sup>th</sup> Percentile	Max Daily Value
		1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>			
Seattle Beacon Hill*	69	4.9	3.7	--	--	--	22.2	103
Seattle Duwamish	43	7.3	5.1	--	14.3	--	34.4	34.4

Notes:

- (1) Sampling occurs for a 24-hour period from midnight to midnight.
- (2) Quarterly averages are shown only if 75 percent or more of the data are available. For both sites Q3 sample percentages were below 75%, and for Seattle Beacon Hill Q4 sample percentages were below 75%.
- (3) Annual averages are shown only if there is at least 75 percent of each of the 4 quarterly averages.
- (4) Data from primary sampler at site

\* Operated by the Washington State Department of Ecology.

## Particulate Matter (PM<sub>2.5</sub>) – Federal Equivalent Methods

Micrograms per Cubic Meter

Equivalent Sampling Methods: Met One BAM

2022

Location	Number of Values	Quarterly Arithmetic Averages				Annual Mean	98th Percentile	Max Daily Value
		1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>			
Auburn	358	8.6	5.1	8.6	13.2	8.9	29.8	86.6
Bremerton Spruce	359	6.2	4.0	5.9	9.4	6.4	19.6	52.5
Darrington	349	10.1	3.4	8.8	26.5	12.2	69.4	255.9
Kent	358	7.9	4.2	9.1	15.8	9.3	33.7	109.5
Marysville	355	9.0	4.2	5.6	17.7	9.1	38.1	114.4
Seattle 10 <sup>th</sup> and Weller*	303	8.1	5.0	9.4	--	--	29.7	108.0
Seattle Beacon Hill*	344	4.9	3.8	7.3	12.1	7.0	27.7	110.4
Seattle Duwamish	360	8.3	5.0	6.8	15.5	8.9	27.7	110.4
Tukwila Allentown	359	6.8	4.1	8.0	13.5	8.1	30.5	99.1
Tacoma South L St	337	8.2	4.5	7.3	13.4	8.3	32.1	89.3
Tacoma South 36 <sup>th</sup> S*	355	8.4	3.6	6.7	16.1	8.7	38.1	94.4
Tacoma Tideflats	344	8.7	4.9	7.2	13.6	8.6	33.6	86.6

Notes:

- (1) Sampling occurs continuously for 24 hours each day.
- (2) Quarterly averages are shown only if 75 percent or more of the data for the quarter is available.
- (3) Annual averages are shown only if 75 percent or more of the data for each of the 4 quarters is available.
- (4) Data from primary sampler at site.

\* Operated by the Washington State Department of Ecology.

## Particulate Matter (PM<sub>2.5</sub>) – Continuous – Nephelometer

Micrograms per Cubic Meter

Sampling Method: Ecotech Nephelometer

2022

Location	Number of Values	Quarterly Arithmetic Averages				Annual Mean	98th Percentile	Max Daily Value
		1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>			
Auburn	325	7.1	4.4	9.2	14.8	8.9	38.5	121
Bellevue**	365	2.7	2	5.2	9.7	4.9	29.5	129.5
Bremerton Spruce*	355	4.8	4.2	7.6	10	6.6	21.6	64.4
Darrington	364	7.8	2.6	8.5	24.5	10.9	76.1	262.7
Kent	362	6.7	4.7	8.5	16.1	9	38.2	146
Lake Forest Park	359	6.6	3.6	7.3	14.1	7.9	33.3	126.9
Marysville	353	7.2	5.3	9.2	19.4	10.3	38.3	174.5
North Bend**	363	2.3	1.8	6.5	11.5	5.5	32	161.4
Seattle Duwamish	347	9.5	5.7	8.5	10.3	8.5	28.6	51.7
Seattle South Park*	364	8.1	6	9.6	14.4	9.5	31.1	106.8
Tukwila Allentown	364	7	4.5	8.4	14.8	8.7	35	140.4
Tacoma South L St	346	8.6	4.7	8.7	17.5	9.9	44.7	134.6
Tacoma Tideflats	357	6.3	3.7	5.5	9.5	6.3	22.6	78.3

Notes:

- (1) Sampling occurs continuously for 24 hours each day.
- (2) Quarterly averages are shown only if 75 percent or more of the data for the quarter is available.
- (3) Annual averages are shown only if 75 percent or more of the data for each of the 4 quarters is available.
- (4) All data values are calculated using site-specific relationships with Federal Reference Method samplers when available.

\*Not available at these sites.

\*\* Operated by the Washington State Department of Ecology.

(5) Data from primary sampler at site.

## **PM<sub>2.5</sub> Speciation Analytes Monitored in 2022**

in Micrograms per Cubic Meter

Acceptable Pm2.5 Aqi & Speciation Mass	Oc Csn_Rev Unadjusted Pm2.5 Lc Tot
Aluminum Pm2.5 Lc	Oc Pm2.5 Lc Tor
Ammonium Ion Pm2.5 Lc	Oc Pm2.5 Lc Tot
Ammonium Nitrate Pm2.5 Lc	Oc1 Csn_Rev Unadjusted Pm2.5 Lc
Ammonium Sulfate Pm2.5 Lc	Oc1 Pm2.5 Lc
Antimony Pm2.5 Lc	Oc2 Csn_Rev Unadjusted Pm2.5 Lc
Arsenic Pm2.5 Lc	Oc2 Pm2.5 Lc
Barium Pm2.5 Lc	Oc3 Csn_Rev Unadjusted Pm2.5 Lc
Bromine Pm2.5 Lc	Oc3 Pm2.5 Lc
Cadmium Pm2.5 Lc	Oc4 Csn_Rev Unadjusted Pm2.5 Lc
Calcium Pm2.5 Lc	Oc4 Pm2.5 Lc
Cerium Pm2.5 Lc	Op Csn_Rev Unadjusted Pm2.5 Lc Tor
Cesium Pm2.5 Lc	Op Csn_Rev Unadjusted Pm2.5 Lc Tot
Chloride Pm2.5 Lc	Op Pm2.5 Lc Tor
Chlorine Pm2.5 Lc	Op Pm2.5 Lc Tot
Chromium Pm2.5 Lc	Organic Carbon Mass Pm2.5 Lc
Cobalt Pm2.5 Lc	Phosphorus Pm2.5 Lc
Copper Pm2.5 Lc	Pm2.5 - Local Conditions
Ec Csn_Rev Pm2.5 Lc Tor	Potassium Ion Pm2.5 Lc
Ec Csn_Rev Pm2.5 Lc Tot	Potassium Pm2.5 Lc
Ec Pm2.5 Lc Tor	Rubidium Pm2.5 Lc
Ec Pm2.5 Lc Tot	Selenium Pm2.5 Lc
EC1 CSN_Rev Unadjusted PM2.5 LC	Silicon Pm2.5 Lc
Ec1 Pm2.5 Lc	Silver Pm2.5 Lc
EC2 CSN_Rev Unadjusted PM2.5 LC	Sodium Ion Pm2.5 Lc
Ec2 Pm2.5 Lc	Sodium Pm2.5 Lc
EC3 CSN_Rev Unadjusted PM2.5 LC	Soil Pm2.5 Lc
Ec3 Pm2.5 Lc	Strontium Pm2.5 Lc
Indium Pm2.5 Lc	Sulfate Pm2.5 Lc
Iron Pm2.5 Lc	Sulfur Pm2.5 Lc
Lead Pm2.5 Lc	Tin Pm2.5 Lc
Magnesium Pm2.5 Lc	Titanium Pm2.5 Lc
Manganese Pm2.5 Lc	Total Nitrate Pm2.5 Lc
Nickel Pm2.5 Lc	Vanadium Pm2.5 Lc
Nitrite Pm2.5 Lc	Zinc Pm2.5 Lc
Oc Csn_Rev Unadjusted Pm2.5 Lc Tor	Zirconium Pm2.5 Lc

Additional information can be obtained at: [aqs.epa.gov/aqsweb/documents/data\\_mart\\_welcome.html](http://aqs.epa.gov/aqsweb/documents/data_mart_welcome.html)

## PM<sub>2.5</sub> Black Carbon

Micrograms per Cubic Meter

Sampling Method: Light Absorption by Aethalometer  
2022

Location	Number of Values	Quarterly Arithmetic Averages				Annual Mean	Max Daily Value
		1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>		
Seattle Duwamish	365	1.0	0.5	0.8	1.3	0.9	4.8
Tukwila Allentown	321	1.1	--	0.5	1.4	--	4.8
Kent	363	0.8	0.4	0.6	1.1	0.7	3.1
Tacoma Tideflats	349	1.2	0.7	0.9	1.5	1.1	5.4
Seattle 10 <sup>th</sup> & Weller	347	1.3	1.0	1.3	1.0	1.2	4.0
Tacoma South L St	363	1.1	0.6	0.8	1.3	0.9	4.2
Tacoma South 36th S	238	0.7	0.2	--	--	--	3.0

Notes:

- (1) Sampling occurs continuously for 24 hours each day.
- (2) Quarterly averages are shown only if 75 % or more of the data is available.
- (3) Annual averages are shown only if there is at least 75 percent of each 4 quarterly averages.

## Ozone (8-hour concentration)

Parts per Million

Sampling Method: Ultraviolet Photometric Detection Method  
2022

Location / Continuous Sampling Period(s)	2021 4 <sup>th</sup> Highest Daily 8-Hour Concentration		4 <sup>th</sup> Highest Daily 8-Hour Concentration			3-Year Average of 4 <sup>th</sup> Highest 8-Hour Concentration
	Value	Date	2020	2021	2022	
Seattle Beacon Hill (1 Jan- 31 Dec)	0.047	29-Apr	0.052	0.052	0.047	0.05
Lake Sammamish Park (1 May – 30 Sep)	0.065	31-Jul	0.06	--	0.065	--
North Bend (1 May – 30 Sep)	0.067	26-Jul	0.051	0.055	0.067	0.057
Enumclaw Mud Mountain (1 May – 30 Sep)	0.075	27-Jun	0.059	0.078	0.075	0.07
Yelm (1 May – 30 Sep)	--	--	0.057	--	--	--
Mt Rainier National Park (1 Jan – 31 Dec)	0.06	8-Aug	0.06	0.058	0.06	0.059

Notes:

- (1) All ozone stations operated by the Washington State Department of Ecology.
- (2) Ending times are reported in Pacific Standard Time.
- (3) For equal concentration values the date and time refer to the earliest occurrences.
- (4) Continuous sampling periods are those with fewer than 10 consecutive days of missing data.

### 2022 Beacon Hill Air Toxics Statistical Summary for Air Toxics (units in parts per billion)

	1,3-Butadiene	Acetaldehyde	Acrolein	Benzene	Carbon Tetrachloride	Chloroform	Ethylbenzene	Ethylene Dichloride	Ethylene Oxide	Formaldehyde	Tetrachloro ethylene
2022 Count	57	60	56	57	57	57	57	57	57	60	57
ND's (reported as 0)	2	0	0	0	0	0	0	0	1	0	0
Median (ppb)	0.0149	0.6830	0.1810	0.1470	0.0804	0.0203	0.0370	0.0147	0.0804	1.0600	0.0082
Mean (ppb)	0.0182	0.9872	0.2071	0.1666	0.0820	0.0214	0.0428	0.0203	0.1028	1.2706	0.0096
95th Percentile (ppb)	0.0495	2.5385	0.3458	0.3184	0.0972	0.0301	0.0820	0.0235	0.2526	2.7100	0.0175
Max (ppb)	0.0715	3.2600	0.8220	0.9090	0.0998	0.0333	0.1610	0.3220	0.3720	3.7700	0.0313
# Below MDL	38	1	24	0	0	0	0	1	6	1	56
% Below MDL	67%	2%	43%	0%	0%	0%	0%	2%	11%	2%	98%

Parameters in gray are over 25% below the method detection limit.

ND = Non-Detects (values reported as zero)

MDL = Method Detection Limit

\* = Kaplan-Meier method used to estimate these means due to a large proportion of results being below the MDL.

### 2022 Beacon Hill Air Toxics Statistical Summary for Air Toxics (units in nanograms per cubic meter)

	Arsenic (PM <sub>10</sub> )	Naphthalene
2022 Count	55	59
ND's (reported as 0)	0	0
Median (ng/m <sup>3</sup> )	0.3440	24.200
Mean (ng/m <sup>3</sup> )	0.4953	33.8763
95th Percentile (ng/m <sup>3</sup> )	1.0560	75.8599
Max ng/m <sup>3</sup> )	2.92	193
# Below MDL	1	0
% Below MDL	2%	0%

## Estimates of Air Toxics Risk

Potential cancer risk is estimated by multiplying the concentration of a pollutant by its unit risk factor (URF), a constant that takes into account its cancer potency. This is shown in the equation below:

$$\text{Potential cancer risk} = \text{ambient concentration } (\mu\text{g}/\text{m}^3) * \text{unit risk factor } (\text{risk}/\mu\text{g}/\text{m}^3)$$

Unit risk factors are often based on epidemiological studies (studies of diseases occurring in human populations) and are also extrapolated from laboratory animal studies. Unit risk factors are typically based on an assumed 70-year (lifetime) exposure interval and are available from multiple sources. In this data summary, cancer risk was estimated using unit risk factors from the Washington State Acceptable Source Impact Levels (ASIL) table.<sup>1</sup> The ASIL values relevant to this summary are in the table below. The two sources from which values in the ASIL table are derived are the U.S. EPA's Integrated Risk Information System<sup>2</sup> (IRIS) and California EPA's Office of Environmental Health and Hazard Assessment<sup>3</sup> (OEHHA). Unit risk factors from both of these sources are derived from extensive reviews of peer-reviewed literature and other datasets. The cancer rating, based on IARC definitions, refers to its "weight of evidence" ranking: 1 = carcinogenic to humans, 2A = probably carcinogenic to humans, 2B = possibly carcinogenic to humans, and 3 = not classifiable as to its carcinogenicity to humans.<sup>4</sup>

### 2022 Air Toxics Unit Risk Factors

AIR TOXIC	WA ASIL 460 UNIT RISK FACTOR RISK/ $\mu\text{g}/\text{m}^3$	CANCER RATING <sup>5</sup>
1,3-Butadiene	$3.0 \times 10^{-5}$	1
Acetaldehyde	$2.7 \times 10^{-6}$	2B
Acrolein	$2.9 \times 10^{-6}$	3
Arsenic	$3.3 \times 10^{-3}$	1
Benzene	$7.7 \times 10^{-6}$	1
Cadmium	$4.2 \times 10^{-3}$	1
Carbon Tetrachloride	$5.9 \times 10^{-6}$	2B
Chloroform	$2.3 \times 10^{-5}$	2B
Chromium (Hexavalent)	$2.5 \times 10^{-1}$	1
Ethylbenzene	$2.5 \times 10^{-6}$	2B
Ethylene Dichloride	$2.6 \times 10^{-5}$	2B
Ethylene Oxide	$5.0 \times 10^{-3}$	1
Formaldehyde	$5.9 \times 10^{-6}$	1
Naphthalene	$3.4 \times 10^{-5}$	2B
Tetrachloroethylene	$6.3 \times 10^{-6}$	2A

<sup>1</sup>Washington State Administrative Code. [apps.leg.wa.gov/wac/default.aspx?cite=173-460-150](http://apps.leg.wa.gov/wac/default.aspx?cite=173-460-150).

<sup>2</sup>Integrated Risk Information System, EPA; [epa.gov/iris/](http://epa.gov/iris/).

<sup>3</sup>California EPA, Consolidated Table of OEHHA/ARB-Approved Risk Assessment Health Values, October 2, 2020; <https://ww2.arb.ca.gov/sites/default/files/classic/toxics/healthval/contable.pdf> .

<sup>4</sup>International Agency for Research on Cancer; <http://monographs.iarc.fr/>.

<sup>5</sup>Ratings per International Agency for Research on Cancer, updated July 2019; <http://monographs.iarc.fr/ENG/Classification/>

**2022 Beacon Hill Potential Cancer Risk Estimates per 1,000,000 – 95<sup>th</sup> Percentile****Percentage of samples greater than cancer screen value**

Air Toxic	Rank	Risk based on 95th percentile concentrations (Washington ASIL)	% of samples > ASIL screen
Ethylene oxide	1	2275	98%
Formaldehyde	2	16	98%
Benzene	3	8	98%
Carbon tetrachloride	4	4	100%
Acetaldehyde	5	7	93%
Chloroform	6	3	100%
Arsenic (PM <sub>10</sub> )	7	4	58%
Acrolein	8	2	61%
1,3-Butadiene	9	3	47%
Naphthalene	10	3	36%
Acrylonitrile	11	3	13%
Cadmium (PM <sub>10</sub> )	12	1	4%

## 2022 Non-cancer Reference Concentrations (RfC) and Hazard Indices >1

Air toxic	Non-cancer RfC (ug/m <sup>3</sup> )	Mean Hazard Index
<b>Acrolein</b>	<b>0.35</b>	<b>1.357</b>
Benzene	3	0.177
Formaldehyde	9	0.141
Manganese (PM <sub>10</sub> )	0.09	0.056
Nickel (PM <sub>2.5</sub> )	0.014	0.055
Nickel (PM <sub>10</sub> )	0.014	0.054
Arsenic (PM <sub>10</sub> )	0.015	0.033
Cadmium (PM <sub>2.5</sub> )	0.020	0.031
Manganese (PM <sub>2.5</sub> )	0.09	0.029
1,3-Butadiene	2	0.020
Carbon tetrachloride	40	0.013
Arsenic (PM <sub>2.5</sub> )	0.015	0.011
Acetaldehyde	140	0.007
Dichloromethane	400	0.004
Toluene	300	0.004
Tetrachloroethylene	35	0.002
Beryllium (PM <sub>10</sub> )	0.007	< 0.001
Chloroform	300	< 0.001
Acrylonitrile	5	< 0.001
Ethylene dichloride	400	< 0.001
Styrene	900	< 0.001
Ethylbenzene	2000	< 0.001
Ethylene dibromide	0.8	< 0.001
Carbon disulfide	800	< 0.001
Trichloroethylene	600	< 0.001
Methyl chloroform	1000	< 0.001
Chlorobenzene	1000	< 0.001
Methyl tert-butyl ether	8000	< 0.001

Reference concentrations are based on chronic Reference Exposure Levels (chRELs) from the California Office of Environmental Health Hazard Assessment (OEHHA)<sup>6</sup>.

Mean hazard index, HI = mean concentration/reference concentration.

Acrolein is the only air toxic that fails the screen with a hazard index greater than 1.

<sup>6</sup> <https://oehha.ca.gov/air/general-info/oehha-acute-8-hour-and-chronic-reference-exposure-level-rel-summary>

## 2000-2022 Air Toxics Trends Statistical Summary

The following table includes the statistical information for the potential cancer risk trends found in the data summary, including if the trend is statistically significant at the 95% confidence level.

Air Toxic	Significance (p-value)	Slope (change in risk per million per year)	y-intercept	Correlation (R <sup>2</sup> )	Number of years (N)
1,3-Butadiene	True (0)	-0.110	3.753	0.653	22
Acetaldehyde	True (0)	-0.097	3.967	0.524	22
Acrolein	False (0.585)	-0.039	2.175	0.172	4
Arsenic PM <sub>10</sub>	True (0.003)	-0.050	3.022	0.405	19
Benzene	True (0)	-0.343	10.23	0.707	22
Cadmium PM <sub>10</sub>	False (0.892)	-0.013	1.343	0.001	18
Carbon Tetrachloride	False (0.092)	-0.023	4.172	0.135	22
Chloroform	True (0)	-0.171	5.680	0.775	22
Chromium VI TSP	True (0.005)	-0.713	16.01	0.754	8
Dichloromethane	False (0.258)	0.001	0.009	0.090	16
Ethylbenzene	False (0.249)	-0.010	0.682	0.094	16
Ethylene Dichloride	False (0.368)	0.037	1.110	0.117	9
Ethylene Oxide	False (0.423)	48.09	-266	0.333	4
Formaldehyde	True (0.022)	-0.306	10.63	0.234	22
Naphthalene	True (0.001)	-0.082	2.889	0.784	9
Nickel PM <sub>10</sub>	True (0)	-0.030	0.857	0.723	18
Tetrachloroethylene	True (0)	-0.040	1.265	0.649	22

## Air Quality Standards and Health Goals

Pollutant [links to historical tables of NAAQS reviews]	Primary/ Secondary	Averaging Time	Level	Form
<a href="#">Carbon Monoxide (CO)</a>	primary	8 hours	9 ppm	Not to be exceeded more than once per year
		1 hour	35 ppm	
<a href="#">Lead (Pb)</a>	primary and secondary	Rolling 3 month average	0.15 $\mu\text{g}/\text{m}^3$ <sup>(1)</sup>	Not to be exceeded
<a href="#">Nitrogen Dioxide (NO<sub>2</sub>)</a>	primary	1 hour	100 ppb	98th percentile of 1-hour daily maximum concentrations, averaged over 3 years
	primary and secondary	1 year	53 ppb <sup>(2)</sup>	Annual Mean
<a href="#">Ozone (O<sub>3</sub>)</a>	primary and secondary	8 hours	0.070 ppm <sup>(3)</sup>	Annual fourth-highest daily maximum 8-hour concentration, averaged over 3 years
<a href="#">Particle Pollution (PM)</a>	PM <sub>2.5</sub>	primary	1 year	12.0 $\mu\text{g}/\text{m}^3$ annual mean, averaged over 3 years
		secondary	1 year	15.0 $\mu\text{g}/\text{m}^3$ annual mean, averaged over 3 years
		primary and secondary	24 hours	35 $\mu\text{g}/\text{m}^3$ 98th percentile, averaged over 3 years
	PM <sub>10</sub>	primary and secondary	24 hours	150 $\mu\text{g}/\text{m}^3$ Not to be exceeded more than once per year on average over 3 years
<a href="#">Sulfur Dioxide (SO<sub>2</sub>)</a>	primary	1 hour	75 ppb <sup>(4)</sup>	99th percentile of 1-hour daily maximum concentrations, averaged over 3 years
	secondary	3 hours	0.5 ppm	Not to be exceeded more than once per year

(1) In areas designated nonattainment for the Pb standards prior to the promulgation of the current (2008) standards, and for which implementation plans to attain or maintain the current (2008) standards have not been submitted and approved, the previous standards (1.5  $\mu\text{g}/\text{m}^3$  as a calendar quarter average) also remain in effect.

(2) The level of the annual NO<sub>2</sub> standard is 0.053 ppm. It is shown here in terms of ppb for the purposes of clearer comparison to the 1-hour standard level.

(3) Final rule signed October 1, 2015, and effective December 28, 2015. The previous (2008) O<sub>3</sub> standards additionally remain in effect in some areas. Revocation of the previous (2008) O<sub>3</sub> standards and transitioning to the current (2015) standards will be addressed in the implementation rule for the current standards.

(4) The previous SO<sub>2</sub> standards (0.14 ppm 24-hour and 0.03 ppm annual) will additionally remain in effect in certain areas: (1) any area for which it is not yet 1 year since the effective date of designation under the current (2010) standards, and (2) any area for which an implementation plan providing for attainment of the current (2010) standard has not been submitted and approved and which is designated nonattainment under the previous SO<sub>2</sub> standards or is not meeting the requirements of a SIP call under the previous SO<sub>2</sub> standards (40 CFR 50.4(3)). A SIP call is an EPA action requiring a state to resubmit all or part of its State Implementation Plan to demonstrate attainment of the required NAAQS.

## National Ambient Air Quality Standards (NAAQS)

The [Clean Air Act](#), which was last amended in 1990, requires EPA to set [National Ambient Air Quality Standards](#) (40 CFR part 50) for pollutants considered harmful to public health and the environment. The Clean Air Act identifies two types of national ambient air quality standards. **Primary standards** provide public health protection, including protecting the health of "sensitive" populations such as asthmatics, children, and the elderly. **Secondary standards** provide public welfare protection, including protection against decreased visibility and damage to animals, crops, vegetation, and buildings.

EPA has set National Ambient Air Quality Standards for six principal pollutants, called "criteria" pollutants (listed below). Units of measure for the standards are parts per million (ppm) by volume, parts per billion (ppb) by volume, and micrograms per cubic meter of air ( $\mu\text{g}/\text{m}^3$ ).

EPA is required to re-visit and update standards every five years, to incorporate the latest health and welfare information.

The state of Washington and the Puget Sound region have adopted these standards. For more information, EPA air quality standards and supporting rationale are available at <https://www.epa.gov/criteria-air-pollutants>. Washington State air quality regulations are available at <https://ecology.wa.gov/Regulations-Permits?topics=27>.<sup>7</sup> The air quality standards that apply to the Puget Sound air shed are summarized below.

Pollutants typically have multiple standards with different averaging times; for example, daily and annual standards. Multiple standards are created and enforced to address health impacts as a result of a shorter, high-level exposure versus longer, low-level exposures. These differences are addressed pollutant-by-pollutant. Additional information is on EPA's website at <https://www.epa.gov/criteria-air-pollutants/naaqs-table>

The Agency has developed an air quality health goal for daily PM<sub>2.5</sub> concentrations. The Agency convened a Particulate Matter Health Committee, comprised of local health professionals, who examined the fine particulate health research.<sup>8</sup> The Health Committee did not consider the federal standard at the time to be protective of human health. In 1999, the Agency adopted a health goal of 25  $\mu\text{g}/\text{m}^3$  for a daily average, more protective than the current federal standard of 35  $\mu\text{g}/\text{m}^3$ . This level is consistent with the American Lung Association's goal and the EPA Clean Air Science Advisory Committee's recommended lower range for the EPA's 2006 ambient air quality standard revision.<sup>9</sup> The Agency did not adopt a separate health goal for the annual average.

---

<sup>7</sup>Washington Administrative Code chapters 173-470, 173-474, and 173-475.

<sup>8</sup>Puget Sound Clean Air Agency. Final Report of the Puget Sound Clean Air Agency PM<sub>2.5</sub> Stakeholder Group; October 1999. Report available on request

<sup>9</sup>EPA Clean Air Science Advisory Committee (CASAC) Particulate Matter (PM) Review Panel; <https://yosemite.epa.gov/sab/SABPEOPLE.NSF/PeopleSearch/60BA5C6D6F54A288852568A900645FE4?OpenDocument>.